

Get Set Go for

**NEET | JEE | BOARDS**

www.mtg.in | December 2019 | Pages 92 | ₹ 40


GEAR UP FOR

**JEE** Main 2020

**CHEMISTRY**

India's #1  
CHEMISTRY MONTHLY FOR  
JEE (Main & Advanced) & NEET

t  day



**RANK  
ENHANCER**

**CONCEPT  
MAP**

BRUSH UP *for* **NEET/JEE**

CLASS  
XI-XII

**mtg**

Trust of more than  
1 Crore Readers  
Since 1982



2019120012754

**CBSE**  
warm-up!

CLASS  
XI-XII

**MONTHLY  
TEST DRIVE**

CLASS  
XI-XII



JEE (Advanced) | JEE (Main) | NEET (UG) 2020  
Give Final Touch To Your Preparation

*with*  
**ALLEN's**  
**MAJOR TEST SERIES**

Test Starting From : 22<sup>nd</sup> Dec. 2019



Pen Paper Mode Test

Computer Based Test

Online Test Series



For more details

visit : [dlp.allen.ac.in](http://dlp.allen.ac.in)

Helpline : 0744-2750275

140+ TEST CENTERS  
FULL SYLLABUS TESTS  
ALL INDIA OPEN TEST

DSAT  
(8 Types of Systematic Analysis Of Each Test)

Corporate Office: "Sankalp", CP-6, Indra Vihar, Kota (Raj.) | Website : [allen.ac.in](http://allen.ac.in) | Helpline : 0744-2757575



|| Renuka Mata Prasanna ||

# MOTEGAONKAR

sir's Student  
**SUBODH RANER**  
Came

**ALL  
INDIA  
RANK 1**  
in Chemistry



**Shivraj R. Motegaonkar**

- ✦ M.Sc. Chemistry - Gold Medalist
- ✦ SET/NET-JRF Qualified
- ✦ DRDO Qualified
- ✦ GATE Qualified
- ✦ BARC Qualified
- ✦ TIFR Qualified



**Chem 175**  
SUBODH RANER  
Total-631

**NEET 2019**  
45 out of 45 Questions  
Came from our  
Study Material as it is

**8**

**NEET 2019**  
Student's Secured above  
**170** Marks out of 180

**141**

Student's  
Secured above  
**150** Marks

**62**

Student's  
Secured above  
**600** Marks

**475**

Student's  
Secured above  
**500** Marks



**GANESH KONDAWAR**

**AIR 10**  
AIIMS

▶ NEET-591 ▶ Chem-135



**PRATHMESH ROKDE**

**AIR 26**  
JIPMER  
AIIMS-41

▶ NEET-675 ▶ Chem-170



**ANANT MALPANI**

**AIR 35**  
AIIMS  
JIPMER-72

▶ NEET-665 ▶ Chem-170

प्रा. मोटेगावकर सरांचे

**RCC**

NEET | AIIMS | JIPMER | IIT-JEE

LATUR BRANCH : 9075 40 2222, 9075 41 2222

NANDED BRANCH : 9075 38 2222, 9075 39 2222



Prof. Shivraj Motegaonkar Official

Follow us : [www.rccpattern.com](http://www.rccpattern.com)




mtg

Success Made Simple




# IF YOU AIM TO BE #1, BE WITH THE #1.

## (IT'S THAT SIMPLE!)

#1 BESTSELLER ON AMAZON



AMONG TOP 10 AMAZON BESTSELLERS



### WHAT MAKES US #1:

- Chapterwise student-friendly synopses for quick-and-easy revision • Topicwise MCQs to check your progress
- NCERT Exemplar MCQs • Assertion & Reason questions for an edge in your AIIMS/JEE Preparation
- HOTS MCQs to boost your concepts • 6 Practice Papers for self-assessment • High Definition (HD) multicolour pages

Call 0124-6601200/1800-10-38 673 or email [info@mtg.in](mailto:info@mtg.in) to learn more






# CRACKING JEE IS THIS SIMPLE



## ALL THE HELP YOU NEED TO CRACK JEE:

- Based on latest pattern of JEE Main • Covers the entire syllabus
- Full of graphic illustration for deep understanding of concepts
- Level wise MCQs with detailed solutions
- NCERT Drill MCQs framed from NCERT Books
- Previous 10 Years' MCQs (2018-2009) of JEE Main/AIEEE
- JEE Main **January & April 2019** Solved Papers Included




Available at all leading book shops throughout India. To buy online visit [www.mtg.in](http://www.mtg.in)

For more information or for help in placing your order, call 0124-6601200 or email [info@mtg.in](mailto:info@mtg.in)



**EARLY** TO  
**RESONANCE,**  
**EARLY** TO  
**SUCCESS.**



# ADMISSION OPEN

**For Classes: V to XII & XII+ (XII Passed)**

**JEE ADVANCED | JEE MAIN | PRE-MEDICAL  
COMMERCE & LAW | PCCP (V to X)  
OLYMPIADS | NTSE | IJSO | KVPY**

**Enroll Now for Academic Session 2020-21  
@ Coaching Fee of 2019-20**

**Call Now: 1800 258 5555**

## Academic Benefits\*

- More than 800 Academic Hours
- More than 500 Classes
- More than 100 Testing Hours

## Financial Benefits\*

- Upto Rs.35000+ Saving on 1 Year Course Fee
- 50% Concession on Admission Form Fee
- Up to 90% Scholarship on Course Fee

**ResoFAST Test Dates: 8<sup>th</sup>, 22<sup>nd</sup> & 29<sup>th</sup> December 2019**

**Resonance Eduventures Limited: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005**

**Tel. No.: 0744-2777777, 2777700 | CIN: U80302RJ2007PLC024029**

**website: [www.resonance.ac.in](http://www.resonance.ac.in) | e-mail: [contact@resonance.ac.in](mailto:contact@resonance.ac.in)**    



# CHEMISTRY today

Volume 28

No. 12

December 2019

**Managing Editor**  
Mahabir Singh

**Editor**  
Anil Ahlawat

**Corporate Office:**

Plot 99, Sector 44 Institutional area, Gurugram -122 003 (HR).  
Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

**Regd. Office:**

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

## CONTENTS

### Class 11

Get Set Go for NEET	8
Brush Up for NEET/JEE	13
CBSE Warm Up	26
Monthly Test Drive	33

### Class 12

Get Set Go for JEE	37
Brush Up for NEET/JEE	41
Concept Map	46
CBSE Warm Up	56
Monthly Test Drive	63

### Competition Edge

Chemistry Musing Problem Set 77	67
Gear Up for JEE Main 2020	69
Concept Booster	74
Rank Enhancer	79
Chemistry Musing Solution Set 76	84

Subscribe online at [www.mtg.in](http://www.mtg.in)

Individual Subscription Rates				Combined Subscription Rates			
	9 months	15 months	27 months		9 months	15 months	27 months
Mathematics Today	300	500	850	PCM	900	1400	2500
Chemistry Today	300	500	850	PCB	900	1400	2500
Physics For You	300	500	850	PCMB	1200	1900	3400
Biology Today	300	500	850				

Send D.D./M.O in favour of MTG Learning Media (P) Ltd.  
Payments should be made directly to : MTG Learning Media (P) Ltd,  
Plot No. 99, Sector 44, Gurugram - 122003 (Haryana)  
We have not appointed any subscription agent.

Printed and Published by Mahabir Singh on behalf of MTG Learning Media Pvt. Ltd. Printed at HT Media Ltd., B-2, Sector-63, Noida, UP-201307 and published at 406, Taj Apartment, Ring Road, Near Safdarjung Hospital, New Delhi - 110029.


Editor : Anil Ahlawat

Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.  
Copyright© MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.



# Last-minute check on your NEET readiness



₹ 625



MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success. **Order now!**



Scan now with your smartphone or tablet\*

## HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 17 years' solved test papers of AIPMT / NEET
- Includes NEET 2019 solved paper
- Detailed solutions for self-assessment and to practice time management



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)

\*Application to read QR codes required

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!



# GET SET GO

# NEET






Exam on  
May 3, 2020



*with exclusive and brain storming MCQs*

Practicing these MCQs help to strengthen your concepts and give you extra edge in your NEET preparation

1. Which of the following is most stable enol form of the given compound?






2. Which of the following pairs can be distinguished by action of heat?

- (i)  $K_2CO_3$  and  $CaCO_3$   
 (ii)  $Na_2CO_3$  and  $Mg(NO_3)_2$   
 (iii)  $Mg(NO_3)_2$  and  $NaNO_3$   
 (a) Only (i) and (ii)      (b) Only (i) and (iii)  
 (c) (i), (ii) and (iii)      (d) None of these


3. Generally commercial hydrochloric acid is prepared by heating NaCl with concentrated  $H_2SO_4$ . How much  $H_2SO_4$  solution containing 93%  $H_2SO_4$  by mass is required for the production 1000 kg of concentrated hydrochloric acid containing 43% HCl by weight?

- (a) 590.0 kg solution of  $H_2SO_4$   
 (b) 310.3 kg solution of  $H_2SO_4$   
 (c) 620.7 kg solution of  $H_2SO_4$   
 (d) 708.2 kg solution of  $H_2SO_4$

4. The major product for the following reaction is







5. 1 mol  $N_2$  and 4 mol  $H_2$  are taken in 15 L flask at  $27^\circ C$ .



After complete conversion of  $N_2$  into  $NH_3$ , 5 L of  $H_2O$  is added. Pressure set up in the flask is

- (a)  $\frac{3 \times 0.0821 \times 300}{15}$  atm (b)  $\frac{2 \times 0.0821 \times 300}{10}$  atm  
(c)  $\frac{1 \times 0.0821 \times 300}{15}$  atm (d)  $\frac{1 \times 0.0821 \times 300}{10}$  atm

6. When  $NaBH_4$  is dissolved in water,  
(a)  $Na^+$  and  $BH_4^-$  ions are formed which are stable.  
(b) it evolve  $H_2$  gas.  
(c)  $NaOH$  and  $B_2O_3$  is formed.  
(d)  $NaOH$  and  $H_3BO_3$  is formed.

7. Alkene (Z)  $\xrightarrow[Zn, H_2O]{O_3}$   $CH_3-\overset{\overset{O}{\parallel}}{C}-Ph +$   
 $CH_3-\overset{\overset{O}{\parallel}}{C}-CH=O + CH_3-CH=O$

Alkene (Z) can be

- (a)  $CH_3-\overset{\overset{O}{\parallel}}{C}-CH=CH-Ph$   
 $CH_3-\overset{\overset{O}{\parallel}}{C}-CH_3$   
(b)  $CH_3-\overset{\overset{O}{\parallel}}{C}-Ph$   
 $CH_3-\overset{\overset{O}{\parallel}}{C}-CH=CH-CH_3$   
(c)  $CH_3-\overset{\overset{O}{\parallel}}{C}=CH-Ph$   
 $CH_3-\overset{\overset{O}{\parallel}}{C}=CH-CH_3$   
(d)  $Ph-\overset{\overset{O}{\parallel}}{C}-CH=CH-CH_3$   
 $CH_3-\overset{\overset{O}{\parallel}}{C}-CH_3$

8. Which of the following transition in hydrogen atom emits photon of maximum frequency?  
(a) Second spectral line of Balmer series  
(b) Second spectral line of Paschen series  
(c) Fifth spectral line of Humphery series  
(d) First spectral line of Lyman series.
9. Which of the following should have pyramidal shape?  
(a)  $[ClOF_2]^+$  (b)  $ICl_3$   
(c)  $[BrICl]^-$  (d) All of these
10. For a hypothetical reversible reaction,  
 $1/2A_{2(g)} + 3/2B_{2(g)} \rightleftharpoons AB_{3(g)}; \Delta H = -20 \text{ kJ}$  if  
standard entropies of  $A_2$ ,  $B_2$  and  $AB_3$  are 60, 40 and

$50 \text{ JK}^{-1} \text{ mole}^{-1}$  respectively. The above reaction will be in equilibrium at

- (a) 400 K (b) 500 K (c) 250 K (d) 200 K
11. Which of the following have been arranged in order of decreasing oxidation number of sulphur?  
(a)  $H_2S_2O_7 > Na_2S_4O_6 > Na_2S_2O_3 > S_8$   
(b)  $SO > SO_4^{2-} > SO_3^{2-} > HSO_4^-$   
(c)  $H_2S > SCl_2 > H_2SO_3 > H_2SO_5$   
(d)  $H_2SO_4 > SO_2 > H_2S > H_2S_2O_8$
12. Which one of the following statements is correct?  
(a) The elements having large negative values of electron gain enthalpy generally act as strong oxidising agents.  
(b) The elements having low values of ionisation enthalpies act as strong reducing agents.  
(c) The formation of  $S_{(g)}^{2-}$  from  $S_{(g)}$  is an endothermic process.  
(d) All of the above.

13. If  $A \rightleftharpoons B (K_c = 3)$ ,  $B \rightleftharpoons C (K_c = 5)$ ,  $C \rightleftharpoons D (K_c = 2)$   
The value of equilibrium constant for the above reaction are given, the value of equilibrium constant for  $D \rightleftharpoons A$  will be  
(a) 15 (b) 0.3 (c) 30 (d) 0.03

14. Which of the following statements is false?  
(a) Absorption of the terrestrially radiated heat by the carbon dioxide is the main cause of global warming.  
(b) Over Antarctica, the depletion of ozone layer is due to the formation of chlorine nitrate.  
(c) The global warming of the earth surface is mainly due to reforestation.  
(d)  $CO_2$ ,  $NO$ ,  $CH_4$ ,  $O_3$ ,  $CCl_4$  and water vapour are greenhouse gases.

15. Metals like Pt and Pd can take up large volume of hydrogen under special conditions. Hydrogen thus retained by the metal is called  
(a) absorbed hydrogen (b) nascent hydrogen  
(c) reactive hydrogen (d) occluded hydrogen.

## SOLUTIONS

1. (c) : Repulsion between two  $C=C$  group of keto form is converted into more stabilized enol form by hydrogen bonding.
2. (c) :  $Na_2CO_3$  and  $K_2CO_3 \xrightarrow{\Delta}$  no reaction  
 $CaCO_3 \xrightarrow{\Delta} CaO + CO_2$




# NEET

Revision cum Crash Course


TEST YOURSELF

PUSH YOUR LIMITS


REACH FOR YOUR PEAK POTENTIAL



₹400/-



₹400/-



₹400/-

## Key Features



- Syllabus systematically divided into 30 days
- NCERT based crisp theory
- Exam Drill exercises for each day with detailed solutions
- OMR sheets at the end of each exercise to improve the skills to attempt actual question paper
- 5 Unit Tests for formative assessment during the preparation
- 2 Mock Test Papers at the end to give the real exam feeling
- NEET 2019 solved paper included

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!

Available at all leading book shops throughout India. To buy online visit [www.mtg.in](http://www.mtg.in).  
For more information or for help in placing your order,  
call 0124-6601200, 1800-10-38673 (Toll Free) or e-mail [info@mtg.in](mailto:info@mtg.in)

Also available on  
Flipkart, Snapdeal, Amazon,  
Infibeam on cash on delivery









$$\text{Mass of HCl} = 1000 \times \left( \frac{43}{100} \right) = 430 \text{ kg}$$

$$\frac{\text{Moles of HCl}}{2} = \frac{\text{Moles of H}_2\text{SO}_4}{1}$$

$$\Rightarrow \frac{430 \times 10^3}{36.5 \times 2} = \text{moles of H}_2\text{SO}_4$$

$$\text{Mass of H}_2\text{SO}_4 = \frac{98 \times 430 \times 10^3}{36.5 \times 2} = 577.26 \times 10^3 \text{ g}$$





$$\text{Mass of 93\% H}_2\text{SO}_4 = 577.26 \times \frac{100}{93} = 620.71 \text{ kg}$$



$$\begin{array}{cccc} t=0 & 1 \text{ mole} & 4 \text{ mole} & 0 \\ t=t_{\text{final}} & 0 & 1 \text{ mole} & 2 \text{ mole} \end{array}$$

$\text{NH}_3$  will be absorbed by water and volume will be  $15 - 5 = 10 \text{ L}$

$$P = \frac{nRT}{V} = \frac{1 \times 0.0821 \times 300}{10} \text{ atm}$$



$$\Delta S^\circ = \Sigma S_P^\circ - \Sigma S_R^\circ = 50 - \left( \frac{1}{2} \times 60 + \frac{3}{2} \times 40 \right)$$

$$\Delta S^\circ = -40 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$\text{We know that, } \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$\text{At equilibrium, } \Delta G^\circ = 0 \quad \therefore \Delta H^\circ = T\Delta S^\circ$$




$$T = \frac{-20 \times 1000}{-40} = 500 \text{ K}$$



12. (d) : (a) The elements having large negative values of electron gain enthalpy generally act as strong oxidising agents *e.g.*, halogens.

(b) The elements having low values of ionisation enthalpies act as strong reducing agents. *e.g.*, alkali metals.

(c) The formation of  $\text{S}_{(g)}^{2-}$  from  $\text{S}_{(g)}$  is an endothermic process as removal of electron will require energy.



$$\text{or } K_c = K_{c_1} \times K_{c_2} \times K_{c_3} = 3 \times 5 \times 2 = 30$$

$$\text{For } D \rightleftharpoons A ; K'_c = \frac{[A]}{[D]} ; K'_c = \frac{1}{K_c} = \frac{1}{30} = 0.03$$



15. (d) : Metal like Pt and Pd, Ni etc. absorb large quantities of the gases, due to vacancies between these atom. This is called occlusion and hydrogen is called occluded hydrogen.

## UNSCRAMBLED WORDS

NOVEMBER 2019

1-c- VERDIGRIS

2-e- TERATOGEN

3-a- AGLYCON

4-g- TRANSFERRIN

5-b- FLOCCULENT

6-d- ELIXIRS

7-h- CLARIFIER

8-f- MUCILAGES



# BRUSH UP for NEET/JEE 2020

Brush up your concepts to get high rank in NEET/JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

## Unit 6

## The *p*-Block Elements

### INTRODUCTION

- Elements having outer electronic configuration of the type  $ns^2 np^1$  to  $ns^2 np^6$ , with all other inner orbitals completely filled, are termed as *p*-block elements. In other words, the elements in which the last electron enters into any of the outermost *p*-orbitals are called *p*-block elements. Thus, the general outer electronic configuration of the *p*-block elements is  $ns^2 np^{1-6}$ .
- The elements belonging to the group 13 to 18 of the long form of periodic table are *p*-block elements. The *p*-block elements include metals, non-metals and metalloids.

### GROUP 13 ELEMENTS : BORON FAMILY

- Group 13 of the periodic table contains six elements, boron (B), aluminium (Al), gallium (Ga), indium (In), thallium (Tl) and Nihonium (Nh).

### Electronic Configuration

- Their valence shell electronic configuration is  $ns^2 np^1$ .

Element	Symbol	Electronic configuration [noble gas] $ns^2 np^1$
Boron	${}_5\text{B}$	$[\text{He}]2s^2 2p^1$
Aluminium	${}_{13}\text{Al}$	$[\text{Ne}]3s^2 3p^1$
Gallium	${}_{31}\text{Ga}$	$[\text{Ar}]3d^{10} 4s^2 4p^1$
Indium	${}_{49}\text{In}$	$[\text{Kr}]4d^{10} 5s^2 5p^1$
Thallium	${}_{81}\text{Tl}$	$[\text{Xe}]4f^{14} 5d^{10} 6s^2 6p^1$
Nihonium	${}_{113}\text{Nh}$	$[\text{Rn}]5f^{14} 6d^{10} 7s^2 7p^1$

### General Physical Properties

#### Atomic radii and ionic radii :

- Group 13 elements and their ions have smaller size than the corresponding group 2 elements because of the increase in nuclear charge.
- Their atomic and ionic radii increase on going down the group with an exception at gallium.

Element	Metallic radius (Å)	Ionic radius	
		$M^{3+}$ (Å)	$M^+$ (Å)
B	0.885	0.27	–
Al	1.43	0.535	–
Ga	1.225	0.620	1.20
In	1.67	0.800	1.40
Tl	1.70	0.885	1.50

#### Melting and boiling points :

- Melting points of group 13 elements do not vary regularly due to structural changes in the elements. The melting point decreases from B to Ga and then increases from Ga to Tl.

Element	Melting point (°C)	Boiling point (°C)
B	2300	2550
Al	660	2467
Ga	30	2240
In	157	2050
Tl	303	1470

- Boron shows a very high melting point because it exists as a giant covalent polymer crystal structure consisting of icosahedral units.



- **Ionisation enthalpy :** The first ionisation enthalpy values of group 13 elements are lower than the corresponding alkaline earth metals, due to the fact that removal of electron is easy. [ $ns^2 np^1$  configuration].

On moving down the group, *I.E.* decreases from B to Al, but the next element Ga has slightly higher ionisation enthalpy than Al due to the poor shielding of intervening *d*-electrons. It again decreases in In and then increases in the last element Tl.

- **Oxidation states :**

Valence electronic configuration  $ns^2 np^1$ , suggests that these elements are expected to show +1 and +3 oxidation states.

Element	Oxidation state
B	+3
Al	+1, +3
Ga	+1, +3
In	+1, +3
Tl	+1, +3

- As we move down in the group, stability of +1 oxidation state increases due to the fact that the two *s*-electrons in the outer shell tend to remain paired and do not participate in bonding (inert pair effect).
- The stability of +3 oxidation state decreases from aluminium to thallium.  
Thus,  $B^{3+} > Al^{3+} > Ga^{3+} > In^{3+} > Tl^{3+}$   
 $Tl^+ > In^+ > Ga^+ > Al^+ > B^+$   
 $Ga^{3+} > Ga^+$  while  $Tl^+ > Tl^{3+}$

- **Electropositive or metallic character :**

- Group 13 elements are less electropositive than alkali metals and alkaline earth metals due to their higher ionisation energies.
- The electropositive character first increases down the group from boron to aluminium and then decreases from aluminium to thallium. Al shows higher metallic character than B due to its larger size and lower ionisation energy. Decrease in the electropositive character beyond Al is due to poor shielding by *d*-electrons, so that the valence electrons cannot be easily removed.
- Electropositive character can also be explained on the basis of their respective electrode potential.
- Because of high ionisation energy and small size, B is considered as non-metal.

- **Electronegativity :**

- Group 13 elements are more electronegative than group 1 elements (alkali metals) and group 2 (alkaline earth metals).
- Electronegativity first decreases from B to Al and then increases due to poor shielding by *d*-electrons.

Element	B	Al	Ga	In	Tl
Pauling's electronegativity	2.0	1.5	1.6	1.7	1.8

## TRENDS IN CHEMICAL PROPERTIES

- **Oxides and hydroxides :**


All the elements of group 13 form oxides with formula  $M_2O_3$  and hydroxides with formula  $M(OH)_3$ . Reactivity of these elements towards oxygen increases down the group.

Oxides	Hydroxides	Nature
$B_2O_3$	$B(OH)_3$	acidic
$Al_2O_3$	$Al(OH)_3$	amphoteric
$Ga_2O_3$	$Ga(OH)_3$	amphoteric
$In_2O_3$	$In(OH)_3$	basic
$Tl_2O_3$	$Tl(OH)_3$	strongly basic


As we moves down the group, the atomic size of elements goes on increasing whereas, the ionisation energy decreases. As a result, the strength of *M* – O bond goes on decreasing accounting for the increase in basic character down the group or conversely explains the decrease in acidic character.

- **Halides :** Group 13 elements form trihalides( $MX_3$ ). Monohalides of thallium are also given as  $TlF$ ,  $TlCl$ ,  $TlBr$  and  $TlI$ .

- Due to small size and high electronegativity, all boron halides are covalent and  $sp^2$  hybridised.
- Boron halides  $BX_3$  are electron deficient compounds having six electrons in the outermost orbit thus function as Lewis acid by accepting a pair of electrons.




The Lewis acid strength follows the order :




This order can be explained on the basis of back bonding or back donation. In  $BF_3$ , there is donation of electron from unutilized  $2p$ -orbitals of fluorine to a vacant  $2p$ -orbital of B.






Thus, B–F bond has some double bond character.

- Lewis acid character of halides of group 13 elements decreases in the order :




#### • Formation of hydrides :

- None of the group 13 elements reacts directly with hydrogen. However a number of hydrides of these elements are known.
- Boron forms a number of stable covalent hydrides with general formula  $\text{B}_n\text{H}_{n+4}$  (nido-boranes e.g.,  $\text{B}_2\text{H}_6$ ) and  $\text{B}_n\text{H}_{n+6}$  (arachno-boranes e.g.,  $\text{B}_4\text{H}_{10}$ ). These are called boranes and are electron deficient compounds.
- The simplest and most important boron hydride is diborane ( $\text{B}_2\text{H}_6$ ).
- Other members of this group also form a stable hydride but they are polymeric in nature e.g.,  $(\text{AlH}_3)_n$ ,  $(\text{GaH}_3)_n$ ,  $(\text{InH}_3)_n$  and contain  $\text{M} - \text{H}_b - \text{M}$  bridge bond. Their stability decreases as we move down the group due to corresponding decrease in strength of  $\text{M} - \text{H}$  bond as the size of atom increases.
- Thallium does not form hydrides.




#### • Formation of carbides :

- Boron and aluminium on heating with carbon at high temperature form carbides.



- Boron carbide is covalent in nature.
- Aluminium carbide is ionic carbide. On hydrolysis it gives methane, hence it is sometimes called as aluminium methanide.



### Anomalous Properties of Boron

- Due to the smallest size, high ionisation energy, absence of vacant  $d$ -orbitals and high electronegativity boron shows anomalous behaviour as compared to other members of the group.

Property	Boron	Other elements of group 13
Metallic behaviour	Non-metal	Metals
Maximum covalency	4	6
Allotropy	Exhibits	Do not exhibit
Oxidation states	Only +3	+1, +3
Compounds	Only covalent	Both ionic and covalent
Halides	Monomeric	Polymeric
Aqueous solution	No ionisation	Form cations
Oxides and hydroxides	Acidic	Amphoteric or basic
Action of non-oxidising acids	No action	React
Combination with metals	Forms boride	Do not combine (form alloy)

### PEEP INTO PREVIOUS YEARS

- Which one of the following elements is unable to form  $\text{MF}_6^{3-}$  ion?  
(a) Ga (b) Al (c) B (d) In  
(NEET 2018)

- Consider the following standard electrode potentials ( $E^\circ$  in volts) in aqueous solution,

Element	$\text{M}^{3+}/\text{M}$	$\text{M}^+/\text{M}$
Al	–1.66	+0.55
Tl	+1.26	–0.34




Based on these data, which of the following statements is correct?

- $\text{Tl}^+$  is more stable than  $\text{Al}^+$ .
  - $\text{Tl}^{3+}$  is more stable than  $\text{Al}^{3+}$ .
  - $\text{Al}^+$  is more stable than  $\text{Al}^{3+}$ .
  - $\text{Tl}^+$  is more stable than  $\text{Al}^{3+}$ .

(JEE Main Online 2017)
- The increasing order of atomic radii of the following group 13 elements is  
(a)  $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$  (b)  $\text{Ga} < \text{Al} < \text{In} < \text{Tl}$   
(c)  $\text{Al} < \text{In} < \text{Ga} < \text{Tl}$  (d)  $\text{Al} < \text{Ga} < \text{Tl} < \text{In}$   
(JEE Advanced 2016)



## SOME IMPORTANT COMPOUNDS OF GROUP 13 ELEMENTS

	Preparation	Properties	Structure
Borax ( $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ )	$\text{Ca}_2\text{B}_6\text{O}_{11} + 2\text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 + 2\text{NaBO}_2 + 2\text{CaCO}_3$ <p style="text-align: center;">Colemanite</p> $4\text{NaBO}_2 + \text{CO}_2 \rightarrow \text{Na}_2\text{B}_4\text{O}_7 + \text{Na}_2\text{CO}_3$ $4\text{H}_3\text{BO}_3 + \text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 + 6\text{H}_2\text{O} + \text{CO}_2$	$\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O} \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 + 10\text{H}_2\text{O}$ $\text{Na}_2\text{B}_4\text{O}_7 \xrightarrow{\Delta} 2\text{NaBO}_2 + \text{B}_2\text{O}_3$ <p style="text-align: center;">Transparent glassy bead</p> $\text{Na}_2\text{B}_4\text{O}_7 + 2\text{NaOH} \rightarrow 4\text{NaBO}_2 + \text{H}_2\text{O}$ $\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{B}_4\text{O}_7$	
	<b>Uses :</b> <ul style="list-style-type: none"> <li>It is used as water softener and cleansing agent.</li> <li>In the laboratory it is used for borax bead test.</li> </ul>		
Orthoboric acid ( $\text{H}_3\text{BO}_3$ )	$\text{Na}_2\text{B}_4\text{O}_7 + 2\text{HCl} + 5\text{H}_2\text{O} \rightarrow 4\text{H}_3\text{BO}_3 + 2\text{NaCl}$ $\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{SO}_4 + 5\text{H}_2\text{O} \rightarrow 4\text{H}_3\text{BO}_3 + \text{Na}_2\text{SO}_4$ $\text{Ca}_2\text{B}_6\text{O}_{11} + 4\text{SO}_2 + 11\text{H}_2\text{O} \rightarrow 2\text{Ca}(\text{HSO}_3)_2 + 6\text{H}_3\text{BO}_3$ $\text{B}_2\text{H}_6 + 6\text{H}_2\text{O} \rightarrow 2\text{H}_3\text{BO}_3 + 6\text{H}_2$ $\text{BN} + 3\text{H}_2\text{O} \rightarrow \text{H}_3\text{BO}_3 + \text{NH}_3$	$\text{H}-\text{OH} + \text{B}(\text{OH})_3 \rightarrow [\text{B}(\text{OH})_4]^- + \text{H}^+ ; \quad \text{p}K_a = 9.25$ $\text{H}_3\text{BO}_3 \xrightarrow{370\text{ K}} \text{HBO}_2 + \text{H}_2\text{O}$ $\text{HBO}_2 \xrightarrow[410\text{ K}]{-\text{H}_2\text{O}} \text{H}_2\text{B}_4\text{O}_7$ $\text{H}_2\text{B}_4\text{O}_7 \xrightarrow{\text{Red heat}} 2\text{B}_2\text{O}_3 + \text{H}_2\text{O}$ $\text{B}(\text{OH})_3 + 3\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{Conc. H}_2\text{SO}_4} \text{B}(\text{OC}_2\text{H}_5)_3 + 3\text{H}_2\text{O}$ <p style="text-align: center;">Triethyl borate</p>	
	<b>Uses :</b> <ul style="list-style-type: none"> <li>It is used in the manufacture of heat resistant borosilicate glass.</li> <li>The aqueous solution of boric acid is used as a mild antiseptic especially as eye wash under the name <i>boric lotion</i>.</li> </ul>		
Diborane ( $\text{B}_2\text{H}_6$ )	$2\text{NaBH}_4 + \text{I}_2 \xrightarrow{\text{Diglyme}} \text{B}_2\text{H}_6 + 2\text{NaI} + \text{H}_2$ $2\text{BF}_3 + 6\text{NaH} \xrightarrow{450\text{ K}} \text{B}_2\text{H}_6 + 6\text{NaF}$ $4\text{BF}_3 \cdot \text{Et}_2\text{O} + 3\text{LiAlH}_4 \xrightarrow[\text{ether}]{\text{Diethyl}} 2\text{B}_2\text{H}_6 + 3\text{LiF} + 3\text{AlF}_3 + 4\text{Et}_2\text{O}$	$\text{B}_2\text{H}_6 + 3\text{O}_2 \rightarrow \text{B}_2\text{O}_3 + 3\text{H}_2\text{O}; \quad \Delta_c H^\circ = -1976\text{ kJ mol}^{-1}$ $\text{B}_2\text{H}_6 + 6\text{H}_2\text{O} \rightarrow 2\text{H}_3\text{BO}_3 + 6\text{H}_2$ $\text{B}_2\text{H}_6 + 6\text{CH}_3\text{OH} \rightarrow 2\text{B}(\text{OCH}_3)_3 + 6\text{H}_2$ $\text{B}_2\text{H}_6 + 2\text{NaH} \xrightarrow[\text{ether}]{\text{Diethyl}} 2\text{NaBH}_4$ $\text{B}_2\text{H}_6 + 2\text{LiH} \xrightarrow[\text{ether}]{\text{Diethyl}} 2\text{LiBH}_4$ $\text{B}_2\text{H}_6 + 2\text{KOH} + 2\text{H}_2\text{O} \rightarrow 2\text{KBO}_2 + 6\text{H}_2$ $\text{B}_2\text{H}_6 + 2\text{NMe}_3 \rightarrow 2\text{BH}_3 \cdot \text{NMe}_3$ $3\text{B}_2\text{H}_6 + 6\text{NH}_3 \xrightarrow{\text{low temp}} 3[\text{BH}_2(\text{NH}_3)_2]^+ [\text{BH}_4]^- \xrightarrow{473\text{ K}} 2\text{B}_3\text{N}_3\text{H}_6 + 12\text{H}_2$ <p style="text-align: center;">Borazine</p>	
	<b>Uses :</b> <ul style="list-style-type: none"> <li>It is used for preparing a number of borohydrides such as <math>\text{LiBH}_4</math>, <math>\text{NaBH}_4</math>, etc.</li> <li>It is used as a reducing agent in organic reactions.</li> </ul>		



Aluminium chloride (AlCl <sub>3</sub> )	$2\text{Al} + 3\text{Cl}_2 \longrightarrow 2\text{AlCl}_3$ $2\text{Al} + 6\text{HCl} \longrightarrow 2\text{AlCl}_3 + 3\text{H}_2$ $\text{Al}_2\text{O}_3 + 3\text{C} + 3\text{Cl}_2 \xrightarrow{100^\circ\text{C}} 2\text{AlCl}_3 + 3\text{CO}$	<p>Anhydrous AlCl<sub>3</sub> fumes in moist air due to hydrolysis and the resulting solution is acidic.</p> $\text{AlCl}_3 + 3\text{H}_2\text{O} \longrightarrow \text{Al(OH)}_3 + 3\text{HCl}$ $\text{AlCl}_3 + 3\text{NH}_4\text{OH} \longrightarrow \text{Al(OH)}_3\downarrow + 3\text{NH}_4\text{Cl}$ <p style="text-align: center;">Gelatinous ppt. (Insoluble in excess NH<sub>4</sub>OH)</p> $\text{AlCl}_3 + 3\text{NaOH} \longrightarrow \text{Al(OH)}_3\downarrow + 3\text{NaCl}$ <p style="text-align: center;">(Soluble in excess NaOH)</p> $\text{Al(OH)}_3 + \text{NaOH} \longrightarrow \text{NaAlO}_2 + 2\text{H}_2\text{O}$ <p style="text-align: center;">(Sodium metaaluminate)</p>	
	<p><b>Uses :</b></p> <ul style="list-style-type: none"> <li>Anhydrous AlCl<sub>3</sub> is used as a catalyst in Friedel-Craft's reaction and in cracking of petroleum.</li> </ul>		

### PEEP INTO PREVIOUS YEARS

- Diborane (B<sub>2</sub>H<sub>6</sub>) reacts independently with O<sub>2</sub> and H<sub>2</sub>O to produce, respectively  
 (a) HBO<sub>2</sub> and H<sub>3</sub>BO<sub>3</sub> (b) B<sub>2</sub>O<sub>3</sub> and H<sub>3</sub>BO<sub>3</sub>  
 (c) B<sub>2</sub>O<sub>3</sub> and [BH<sub>4</sub>]<sup>-</sup> (d) H<sub>3</sub>BO<sub>3</sub> and B<sub>2</sub>O<sub>3</sub>  
 (JEE Main 2019)
- The green colour produced in the borax bead test of a chromium(III) salt is due to  
 (a) CrB (b) Cr<sub>2</sub>O<sub>3</sub>  
 (c) Cr(BO<sub>2</sub>)<sub>3</sub> (d) Cr<sub>2</sub>(B<sub>4</sub>O<sub>7</sub>)<sub>3</sub>  
 (JEE Advanced 2019)
- Boric acid is an acid because its molecule  
 (a) contains replaceable H<sup>+</sup> ion  
 (b) gives up a proton  
 (c) accepts OH<sup>-</sup> from water releasing proton  
 (d) combines with proton from water molecule.  
 (NEET Phase-II 2016)

### POINTS FOR EXTRA SCORING

- Gallium is a low melting solid (m.pt = 29.8°C) and readily supercools *i.e.*, remains liquid even at temperatures several degrees below its melting point.
- Boron nitride has a layer structure resembling graphite, in which hexagonal ring consists of alternating of boron and nitrogen atoms.
- Corundum (an α-form of Al<sub>2</sub>O<sub>3</sub>) an extremely hard substance is used as jeweller's rouge to polish glass.

- Boron halides do not exist as dimer due to small size of boron atom which makes it unable to coordinate four large-sized halide ions.
- In thallium triiodide (TlI<sub>3</sub>), thallium is in (+1) oxidation state and anion is I<sub>3</sub><sup>-</sup>.
- Thallium is the highly toxic element amongst the group 13 members.
- The anhydrous alum is called burnt alum.

## GROUP 14 ELEMENTS : CARBON FAMILY

- Group 14 of the periodic table contains six elements, carbon (C), silicon (Si), germanium (Ge), tin (Sn), lead (Pb) and Flerovium (Fl).

### Electronic Configuration

- Their valence shell electronic configuration is  $ns^2np^2$ .

Element	Symbol	Electronic configuration [noble gas] $ns^2np^2$
Carbon	<sub>6</sub> C	[He] $2s^22p^2$
Silicon	<sub>14</sub> Si	[Ne] $3s^23p^2$
Germanium	<sub>32</sub> Ge	[Ar] $3d^{10}4s^24p^2$
Tin	<sub>50</sub> Sn	[Kr] $4d^{10}5s^25p^2$
Lead	<sub>82</sub> Pb	[Xe] $4f^{14}5d^{10}6s^26p^2$
Flerovium	<sub>114</sub> Fl	[Rn] $5f^{14}6d^{10}7s^27p^2$

### General Physical Properties

- Atomic radii :**
  - The atomic radii of group 14 elements are smaller than those of corresponding elements



of group 13, because as we move from left to right in a period, effective nuclear charge increases, consequently, the size of the atom decreases.

- The atomic radii of group 14 elements increase regularly on moving down the group mainly due to increase in the number of shells.

- **Melting and boiling points :**

- The melting and boiling points of group 14 elements are much higher than those of the elements of group 13. It is because these form four covalent bonds with each other and hence there exists strong binding forces between their atoms both in the solid as well as in the liquid state.
- The melting and boiling points decrease on descending the group from carbon to lead because the  $M-M$  bond becomes weaker as the atom size increases.

- **Ionization enthalpy :**

- The first ionization potential of group 14 elements are higher than those of corresponding elements of group 13 due to increased nuclear charge.
- Due to increase in the atomic radii of these elements from C to Pb the ionisation enthalpy decreases on moving down the group.
- Due to ineffective shielding of valence electrons by  $d$ - and  $f$ -orbitals the sequence for ionization energy is




- **Oxidation state :**

Element	Oxidation state
C	+4
Si	+4
Ge	+2, +4
Sn	+2, +4
Pb	+2, +4

The stability of +2 state increases from Ge to Pb, i.e.,  $Ge^{2+} < Sn^{2+} < Pb^{2+}$  but for +4 oxidation state it is  $Ge^{4+} > Sn^{4+} > Pb^{4+}$ .

As we move down, inert pair effect increases and stability of lower oxidation state increases.



- **Electropositive character or metallic character :**

Group 14 elements are less electropositive than group 13 elements.

Element	Metallic character
C	Non-metal
Si	Non-metal
Ge	Metalloid
Sn	Metal
Pb	Metal

- **Catenation (Tendency to form chains) :**

- Catenation is the property of elements to form long chains or rings by self-linking of their own atoms through covalent bonds.
- The tendency for catenation decreases in the order  $C \gg Si > Ge \approx Sn \gg Pb$
- Carbon shows a remarkable catenation due to smaller size and high electronegativity, so that C–C bond is extremely stable.

- **Allotropy :** All the elements of group 14 except Pb show allotropy.

## CHEMICAL PROPERTIES


- **Hydrides :** All members of the group form covalent hydrides. Their number and ease of formation decreases down the group. Hydrides of carbon are called *hydrocarbons* (alkanes, alkenes or alkynes). Hydrides of Si and Ge are known as *silanes* and *germanes*. The only hydrides of Sn and Pb are  $SnH_4$  (stannane) and  $PbH_4$  (plumbane).

- Their thermal stability decreases down the group.
- Their reducing character increases down the group.

- **Halides :** All the elements give tetrahedral and covalent halides of the type  $MX_4$  except  $PbBr_4$  and  $PbI_4$ .

Thermal stability :  $CX_4 > SiX_4 > GeX_4 > SnX_4 > PbX_4$

Order of thermal stability with common metals :




Except  $CX_4$  other tetrahalides can be hydrolysed due to the presence of vacant  $d$ -orbitals.



Ease of hydrolysis :  $SiX_4 > GeX_4 > SnX_4 > PbX_4$

- **Oxides :** They form two type of oxides, monoxides of the type  $MO$  and dioxides of the type  $MO_2$ .



$SiO_2$  is a solid with three dimensional network in which Si is bonded to four oxygen atoms tetrahedrally and covalently.



- **Carbides :**

- The binary compounds of carbon with elements other than hydrogen are called *carbides*.
- Ionic carbides are formed by the most electropositive metals such as alkali and alkaline earth metals and Al.
- Both  $\text{Be}_2\text{C}$  and  $\text{Al}_4\text{C}_3$  are called methanides because they react with  $\text{H}_2\text{O}$  yielding methane.
- Covalent carbides are formed by Si and B.
- SiC (carborundum) has a diamond like structure, hence it is called *artificial diamond*.
- Interstitial carbides are formed by transition elements in which C-atoms occupy interstitial positions in the close-packed arrays of metal atoms. W, Zr, Ti, Ta and Mo can form interstitial carbides.

## Allotropes of Carbon


- The allotropic forms of carbon are :
  - Crystalline form : diamond and graphite.
  - Amorphous form : coal, charcoal, lampblack, etc.

### Diamond :

- It is a transparent, crystalline substance with very high refractive index.
- It is the purest form of carbon found naturally and can also be made artificially.
- Diamond is the hardest natural substance known and is a bad conductor of electricity.

### Structure :

- Each carbon atom of diamond is bonded to four other carbon atoms, through  $sp^3$  hybridised orbitals, situated at the corners of a regular tetrahedron, with C – C bond length of  $1.54 \text{ \AA}$  and bond angle of  $109.5^\circ$ .
- This gives diamond a three-dimensional network structure.




- **Graphite :**

- It is a dark grey, crystalline solid which is soft and greasy to touch. It possesses a metallic lustre.
- It is also known as '*plumbago*' (black lead) as it leaves a black mark on paper.
- It is a good conductor of electricity and its conductivity increases with rise in temperature.

- **Structure :**


- It has a two dimensional sheet like structure with each carbon atom being covalently bonded to three carbon atoms through  $sp^2$  hybridised orbitals, forming a planar hexagonal structure.
- The fourth electron of each carbon forms a  $\pi(\pi)$  bond with partial overlap with that of the neighbouring carbon.
- The C – C bond length is  $1.42 \text{ \AA}$ , shorter than that in diamond. The  $\pi$ -electrons are free to move (mobile electrons) and account for the electrical conductivity.
- The adjacent hexagonal layers (sheets) are held by weak van der Waals' forces thus, making it easy for the layers to slide over one another, accounting for its soft and greasy texture.



### Fullerenes

- Fullerenes are made by heating graphite in an electric arc in the presence of inert gases such as helium or argon. Fullerenes are the only pure form of carbon because they have smooth structure without having 'dangling' bonds. Fullerenes are cage like molecules.  $\text{C}_{60}$  molecule has a shape like soccer ball and called Buckminsterfullerene.
- It contains twenty, six-membered rings and twelve, five-membered rings. A six membered ring is fused with six or five membered ring but a five membered ring can only be fused with six membered ring. All the carbon atoms are equal and they undergo  $sp^2$  hybridisation. This ball shaped molecule has 60 vertices and each one is occupied by one carbon atom and it also contains both single and double bonds with C–C distances of  $143.5 \text{ pm}$  and  $138.3 \text{ pm}$ , respectively. Spherical fullerenes are also called *bucky balls* in short.





Structure of Buckminsterfullerene

## Anomalous Properties of Carbon

- Carbon differs from other elements of group 14 because of its smaller size, high electronegativity, absence of  $d$ -orbitals and high tendency of catenation.

Property	Carbon	Other elements
Hardness	hardest (diamond)	less hard
M.pt. and B.pt.	high	low
Maximum covalency	4	6
Multiple bonds	$p\pi-p\pi$ (high extent)	$p\pi-d\pi$ (low extent)
Catenation	very high tendency	very low
Tetrahalides	does not undergo hydrolysis	undergo hydrolysis

## PEEP INTO PREVIOUS YEARS

7. Which of the following is incorrect statement?  
 (a)  $\text{SnF}_4$  is ionic in nature.  
 (b)  $\text{PbF}_4$  is covalent in nature.  
 (c)  $\text{SiCl}_4$  is easily hydrolysed.  
 (d)  $\text{GeX}_4$  ( $X = \text{F, Cl, Br, I}$ ) is more stable than  $\text{GeX}_2$ .  
 (NEET 2019)
8. A tin chloride 'Q' undergoes the following reactions (not balanced)  
 $\text{Q} + \text{Cl}^- \rightarrow \text{X}$   
 $\text{Q} + \text{Me}_3\text{N} \rightarrow \text{Y}$   
 $\text{Q} + \text{CuCl}_2 \rightarrow \text{Z} + \text{CuCl}$   
 X is monoanion having pyramidal geometry. Both Y and Z are neutral compounds. Choose the correct option(s).  
 (a) The central atom in X is  $sp^3$  hybridized.  
 (b) There is a coordinate bond in Y.  
 (c) The oxidation state of the central atom in Z is +2.  
 (d) The central atom in Z has one lone pair of electrons.  
 (JEE Advanced 2019)
9. In graphite and diamond, the percentage of  $p$ -characters of the hybrid orbitals in hybridisation are respectively  
 (a) 33 and 75  
 (b) 50 and 75  
 (c) 33 and 25  
 (d) 67 and 75  
 (JEE Main Online 2018)



## SOME IMPORTANT COMPOUNDS OF GROUP 14 ELEMENTS

Compound	Preparation	Properties	Uses
<b>Carbon monoxide (CO)</b> $\text{:C}\equiv\text{O:}^+ \leftrightarrow \text{:C}=\ddot{\text{O}}:$ or $\text{:C}\equiv\text{O:}$	Incomplete combustion of carbon and carbon containing compounds. $2\text{C} + \text{O}_2 \rightarrow 2\text{CO}$ Commercially CO is obtained as water gas ( $\text{H}_2 + \text{CO}$ ) and producer gas ( $\text{CO} + \text{N}_2$ ). $\text{C}_{(s)} + \text{H}_2\text{O}_{(g)} \xrightarrow{200-1000^\circ\text{C}} \underbrace{\text{CO}_{(g)} + \text{H}_{2(g)}}_{\text{Water gas}}$	It is neutral towards litmus. It is colourless odourless and tasteless neutral oxide. <div style="text-align: center;"> <math>\begin{array}{c} \text{CO} \\ \swarrow \quad \downarrow \quad \searrow \\ \begin{array}{l} \text{Cl}_2 \rightarrow \text{COCl}_2 \text{ (Phosgene)} \\ \text{Fe}_3\text{O}_4 \xrightarrow{1073\text{ K}} 2\text{Fe} + 3\text{CO}_2 \\ 2[\text{Ag}(\text{NH}_3)_2]^+ \text{OH}^- \rightarrow 2\text{Ag}\downarrow + \text{CO}_2 + \text{H}_2\text{O} + 4\text{NH}_3 \\ \text{ZnO} \xrightarrow{2\text{H}_2} \text{Zn} + \text{CO}_2 \\ \text{Ni} \xrightarrow{323\text{ K}-353\text{ K}} \text{Ni(CO)}_4 \\ \text{CH}_3\text{OH} \end{array} \end{array}</math> </div>	In the form of producer gas ( $\text{N}_2 + \text{CO}$ ) and water gas ( $\text{H}_2 + \text{CO}$ ), it is used as fuel. In the metallurgy of nickel. As a reducing agent. For the manufacture of methyl alcohol and synthetic petrol



	$2C_{(s)} \xrightarrow[\text{Air}]{\begin{matrix} 2g & N_2g \\ \hline & 1000^\circ C \end{matrix}} \xrightarrow[\text{Producer gas}]{\begin{matrix} g & g \end{matrix}}$ $Fe_2O_3 + 3C \xrightarrow{\Delta} 2Fe + 3CO_2$ $ZnO + CO \xrightarrow{\Delta} Zn + CO_2$ $HCOOH \xrightarrow[373\text{ K}]{\text{Conc. } H_2SO_4} CO + H_2O$ $\begin{matrix} COOH \\   \\ COOH \end{matrix} + H_2SO_4 \xrightarrow{\text{Heat}} CO_2 + CO + H_2O$		
<p><b>Silicones (Si—O—Si) linkage <math>(R_2SiO)_n</math></b></p> $\begin{array}{c} R & R \\   &   \\ -O-Si-O-Si-R- \\   &   \\ R & R \end{array}_n$	$2RMgCl + SiCl_4 \longrightarrow R_2SiCl_2 + 2MgCl_2$ <p style="text-align: center;">Dialkyldichloro silane</p> $R_2SiCl_2 \xrightarrow[-2HCl]{+2H_2O} R_2Si(OH)_2$ <p style="text-align: center;">Dialkyldihydro silane</p> $\xrightarrow[-H_2O]{\text{Polymerisation}}$ $\left[ \begin{array}{c} R & R \\   &   \\ -O-Si-O-Si-O- \\   &   \\ R & R \end{array} \right]_n$ <p style="text-align: center;">Silicone</p>	<p>Silicone polymers are highly stable towards heat. High dielectric strength. Polymers of silicon with low molecular weight are soluble in organic solvents.</p> <p>They are not affected by weak acids, alkalies, salt solutions and resistant to oxidation.</p> <p>They are water repellents and good electrical insulators.</p>	<p>Silicone polymers are used as sealant, greases, varnishes and these can be used even at very low temperatures (of the order of <math>-40^\circ C</math>).</p> <p>These are used as lubricants at both high and low temperatures.</p> <p>Silicone rubbers are very useful because they retain their elasticity at lower temperatures as compared to other rubbers. They are also mixed in paints to make them damp resistant.</p>
<p><b>Carbon dioxide (<math>CO_2</math>)</b></p> $\begin{array}{c} \curvearrowright & \curvearrowright \\ :O=C=O: & \longleftrightarrow & ^+O \equiv C-O: \\ & & \longleftrightarrow & ^-O-C \equiv O^+ \end{array}$	$C_{(s)} + O_{2(g)} \longrightarrow CO_{2(g)}$ $C_3H_{8(s)} + 5O_{2(g)} \longrightarrow 3CO_{2(g)} + 4H_2O_{(g)}$ $ZnCO_3 \xrightarrow{\Delta} ZnO + CO_2$ $CaCO_3 \xrightarrow{\Delta} CaO + CO_2$ $NaHCO_3 \xrightarrow{\Delta} Na_2CO_3 + CO_2 \uparrow + H_2O$		<p>In aerated waters, e.g., in soda water, etc.</p> <p><math>CO_2</math> is used in extinguishing fire.</p> <p>Solid carbon dioxide (dry ice) is used as refrigerant.</p> <p>In the manufacture of sodium carbonate and sodium bicarbonate by the Solvay's process.</p>



<p><b>Silicon dioxide or silica (SiO<sub>2</sub>)</b></p>  <p>Three-dimensional structure of SiO<sub>2</sub></p>	$\text{Si} + \text{O}_2 \longrightarrow \text{SiO}_2$ $\text{SiCl}_4 + 2\text{H}_2\text{O} \longrightarrow \text{SiO}_2 + 4\text{HCl}$ $\text{SiF}_4 + 2\text{H}_2\text{O} \longrightarrow \text{SiO}_2 + 4\text{HF}$		<p>As quartz, it is used as a piezoelectric material. In the form of silica gel, it act as drying agent. It also supports chromatographic materials and catalysts.</p>
---	---	---	--

### PEEP INTO PREVIOUS YEARS

10. Which of the following compounds is used in cosmetic surgery?

- (a) Silica (b) Silicates  
(c) Silicones (d) Zeolites

(Odisha NEET 2019)

11. Lithium aluminium hydride reacts with silicon tetrachloride to form

- (a) LiCl, AlCl<sub>3</sub> and SiH<sub>4</sub>  
(b) LiCl, AlH<sub>3</sub> and SiH<sub>4</sub>  
(c) LiH, AlCl<sub>3</sub> and SiCl<sub>2</sub>  
(d) LiH, AlH<sub>3</sub> and SiH<sub>4</sub>

(JEE Main Online 2018)

12. Under hydrolytic conditions, the compounds used for preparation of linear polymer and for chain termination, respectively, are

- (a) CH<sub>3</sub>SiCl<sub>3</sub> and Si(CH<sub>3</sub>)<sub>4</sub>  
(b) (CH<sub>3</sub>)<sub>2</sub>SiCl<sub>2</sub> and (CH<sub>3</sub>)<sub>3</sub>SiCl  
(c) (CH<sub>3</sub>)<sub>2</sub>SiCl<sub>2</sub> and CH<sub>3</sub>SiCl<sub>3</sub>  
(d) SiCl<sub>4</sub> and (CH<sub>3</sub>)<sub>3</sub>SiCl

(JEE Advanced 2015)

### POINTS FOR EXTRA SCORING

- Graphite can be converted into diamond at  $\approx 55 \times 10^3$  atmospheric pressure.
- Chemical composition of ordinary glass is Na<sub>2</sub>SiO<sub>3</sub> · CaSiO<sub>3</sub> · 4SiO<sub>2</sub>.
- PbO is called litharge, Red lead (Pb<sub>3</sub>O<sub>4</sub>) is a mixed oxide.

### Answer Key For Peep Into Previous Years

- |        |          |        |         |         |         |
|--------|----------|--------|---------|---------|---------|
| 1. (c) | 2. (a)   | 3. (b) | 4. (b)  | 5. (c)  | 6. (c)  |
| 7. (b) | 8. (a,b) | 9. (d) | 10. (c) | 11. (a) | 12. (b) |



# WRAP it up!

1. AlCl<sub>3</sub> is an electron deficient compound but AlF<sub>3</sub> is not. This is because

- (a) atomic size of F is smaller than Cl which makes AlF<sub>3</sub> more covalent  
(b) AlCl<sub>3</sub> is a covalent compound while AlF<sub>3</sub> is an ionic compound  
(c) AlCl<sub>3</sub> exists as a dimer but AlF<sub>3</sub> does not  
(d) Al in AlCl<sub>3</sub> is in sp<sup>3</sup> hybrid state but Al in AlF<sub>3</sub> is in sp<sup>2</sup> state.

2. Aqueous solution of borax acts as a buffer because

(a) it contains weak acid and its salt with strong base

- (b) it contains tribasic acid and strong base  
(c) it contains number of neutral water molecules  
(d) none of these.

3. Borosilicate glass has the special property of being

(a) transparent (b) uncrackable  
(c) resistant of heat (d) tough

4. Which of the following exists in gaseous form in nature?

- (a) ClF<sub>3</sub> (b) BF<sub>3</sub>  
(c) IF<sub>3</sub> (d) ICl



5. Pb + conc.  $\text{HNO}_3$  gives  
 (a)  $\text{Pb}(\text{NO}_3)_2 + \text{NO}_2$  (b)  $\text{PbNO}_3 + \text{NO}$   
 (c)  $\text{Pb}(\text{NO}_3)_4 + \text{NO}_3$  (d)  $\text{Pb}(\text{NO}_3)_3 + \text{N}_2\text{O}$
6. A solid element (symbol Y) conducts electricity and forms two chlorides  $\text{YCl}_n$  (a colourless volatile liquid) and  $\text{YCl}_{n-2}$  (a colourless solid). To which one of the following groups of the periodic table does Y belong?  
 (a) 13 (b) 14 (c) 15 (d) 16
7. An element of group 14 forms two oxides one of which is highly poisonous and neutral. Other oxide can be easily liquefied and compressed to give a solid which is used as a refrigerant under the name of drikold. The element and the oxides are  
 (a) Si,  $\text{SiO}$ ,  $\text{SiO}_2$  (b) Pb,  $\text{PbO}$ ,  $\text{PbO}_2$   
 (c) C,  $\text{CO}$ ,  $\text{CO}_2$  (d) Sn,  $\text{SnO}$ ,  $\text{SnO}_2$
8. Which of the following statement about group IV elements (C to Pb) is not correct?  
 (a) Sn and Pb form dichlorides  
 (b) The thermal stability of hydrides decreases from  $\text{CH}_4$  to  $\text{PbH}_4$  as bond enthalpy for  $\text{M}-\text{H}$  bond decreases.  
 (c) All form tetrafluorides, tetrachlorides.  
 (d) All form tetraiodides.
9. Which one of the following has the highest Lewis acid strength?  
 (a)  $\text{BI}_3$  (b)  $\text{BBr}_3$  (c)  $\text{BF}_3$  (d)  $\text{BCl}_3$
10. When bauxite powder is mixed with coke and reacted with nitrogen at 2075 K, carbon monoxide and X are formed. What is the gas formed when X is reacted with water?  
 (a)  $\text{NH}_3$  (b)  $\text{N}_2$  (c)  $\text{N}_2\text{O}$  (d)  $\text{O}_2$
11. Which of the following is strongest bond?  
 (a)  $\text{Si}-\text{Si}$  (b)  $\text{C}-\text{C}$   
 (c)  $\text{Sn}-\text{Sn}$  (d)  $\text{Pb}-\text{Pb}$
12. When  $\text{NaOH}$  solution is added to  $\text{AlCl}_3$  solution, a white ppt. first appears which dissolves in excess of  $\text{NaOH}$ . In the solution so obtained  
 (a) aluminium ion is present in anion  
 (b) aluminium ion is present in cation  
 (c) aluminium ion is present in both the ions  
 (d) aluminium is present in neither the cation nor the anion.
13. Iodine oxidises sodium borohydride to give  
 (a)  $\text{B}_2\text{H}_6$  (b) sodium hydride  
 (c)  $\text{HI}$  (d)  $\text{I}_3^-$
14. In the manufacture of cement which of the following is used?

## UNSCRAMBLE ME

Unscramble the words given in column I and match them with their explanations in column II.

### Column I

1. TESOROEC
2. PASTSELATHI
3. PXECILEX
4. TAGCYUFI
5. LMULIET
6. ROXATOHNL
7. NATPAI
8. RAPSILECE

### Column II

- (a) A molecular charge transfer complex formed by the complexing of a donor in an excited state with an acceptor in its ground state.
- (b) Used in the measure of effective pressure dependence versus the chemical potential of a gas.
- (c) Fractions obtained in the distillation of coal or wood tar.
- (d) The decorative and corrosion-resistant surface given to bronze, iron, etc., as an oxide film formed on warming the metal.
- (e)  $\text{MgO}$  either natural or formed by heating  $\text{Mg}(\text{OH})_2$  used in furnace linings.
- (f) Naturally occurring, hard bitumens of high purity containing only 1-2% of mineral matter.
- (g) Reduction product of anthraquinone.
- (h)  $\text{Al}_6\text{Si}_2\text{O}_{13}$ , the stablest aluminum silicate, formed by heating other aluminosilicates used in refractories and glasses.

Readers can send their responses at [editor@mtg.in](mailto:editor@mtg.in) or post us with complete address by 10<sup>th</sup> of every month.  
 Names of solution senders will be published in next issue.




- (a) Clay and silica (b) Limestone and silica  
(c) Limestone and clay (d) Limestone and gypsum.

15. Which is not correct?

- (a)  $\text{GeO}_2$  is weakly acidic.  
(b)  $\text{GeCl}_4$  forms  $[\text{GeCl}_2]^{2-}$  ion with  $\text{HCl}$ .  
(c)  $\text{Ge}(\text{OH})_2$  is amphoteric.  
(d)  $\text{GeCl}_2$  is more stable than  $\text{GeCl}_4$ .

16. The products of the following reaction are



- (a)  $\text{SiC}$  and  $\text{CO}_2$  (b)  $\text{SiO}$  and  $\text{CO}$   
(c)  $\text{SiC}$  and  $\text{CO}$  (d)  $\text{Si}$  and  $\text{CO}_2$

17. Which of the following statements is not true both for B and Al?

- (a) They burn in oxygen to give oxides at high temperature.  
(b) Their halides are Lewis acids.  
(c) They combine with nitrogen to form nitrides.  
(d) They react with  $\text{HCl}$  to form chlorides.

18. Unlike the other elements of its group, carbon does not form  $\text{CX}_2$  type molecules because

- (a) energetically this is not possible  
(b) carbon undergoes catenation  
(c) it is non-metallic  
(d) carbon does not contain  $d$ -orbital.

19. Which of the following statements is not correct?

- (a) Silicon is extensively used as a semiconductor.  
(b) Carborundum is  $\text{SiC}$ .  
(c) Silicon occurs in free state in nature.  
(d) Mica contains the element silicon.

20. Covalency of B in  $\text{BF}_4^-$  is

- (a) 5 (b) 4 (c) 3 (d) 2

### SOLUTIONS


1. (b): Due to larger size of Cl,  $\text{AlCl}_3$  is a covalent compound (Fajan's rule).

2. (a)

3. (c): Borosilicate glass is heat resistant, because it has low coefficient of expansion.

4. (b)

5. (a):  $\text{HNO}_3$  is the best solvent for lead.




6. (b): Elements Y can probably be tin which conducts electricity and belongs to group 14.  $\text{SnCl}_4$  is colourless volatile liquid whereas  $\text{SnCl}_2$  is colourless solid.

7. (c): C forms two oxides  $\text{CO}$  and  $\text{CO}_2$ .  $\text{CO}$  is neutral and poisonous.  $\text{CO}_2$  can be compressed to a refrigerant dry ice or solid  $\text{CO}_2$ .  $\text{SiO}$  is unstable.  $\text{SnO}$  and  $\text{PbO}$  are amphoteric in nature.


8. (d): For example,  $\text{PbI}_4$  does not exist because  $\text{Pb}^{4+}$  ion is strong oxidising agent and  $\text{I}^-$  ion is strong reducing agent. Thus,  $\text{Pb}^{4+}$  cannot exist in presence of  $\text{I}^-$  ion and get reduced to  $\text{Pb}^{2+}$  ion.

9. (a)

10. (a): Bauxite powder on heating with coke in presence of  $\text{N}_2$  produces aluminium nitride which on reaction with water produces ammonia.




Bauxite



11. (b): Bond strength of  $M - M$  bond and the tendency for catenation decreases in the order  $\text{C} - \text{C} > \text{Si} - \text{Si} > \text{Ge} - \text{Ge} > \text{Sn} - \text{Sn}$ . Greater the bond energy greater is the strength of bond.


12. (a):  $\text{Al}(\text{OH})_3$  ppt. first formed redissolves in  $\text{NaOH}$  to form sod. meta aluminate where Al is a part of anion  $\text{AlO}_2^-$ .

13. (a): The oxidation of sodium borohydride with iodine in diglyme gives diborane.




14. (b): Limestone and silica are the main components of cement.

15. (d): Ge has greater tendency to show +4 oxidation state.



17. (d): Boron being non-metal does not displace hydrogen from  $\text{HCl}$  to produce its chloride.

18. (a): The elements of group 14 show tetravalency by sharing four of its valence electrons. They have O.S. of +4. Ge, Sn and Pb also show +2 O.S. The stability of dihalides increases down the group. The stability of dihalides follow the sequence



19. (c): Si widely occurs as  $\text{SiO}_2$ .



20. (b): Covalency of B in  $\text{BF}_4^-$  is 4 which is its maximum covalency also.

### EXAM ALERT 2020

Exam	Date
JEE Main I	6 <sup>th</sup> to 11 <sup>th</sup> January
WB JEE	2 <sup>nd</sup> February
JEE Main II	3 <sup>rd</sup> to 9 <sup>th</sup> April
NEET	3 <sup>rd</sup> May
JEE Advanced	17 <sup>th</sup> May



# BEST TOOLS FOR SUCCESS IN **JEE Main**



**10** Very Similar Practice Test Papers with Detailed Solutions

**18** JEE MAIN 2019-2015(Offline & Online)-2013 & AIEEE (2012-2002)  
Years



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!





# CBSE

## warm-up!

CLASS-XI

Chapterwise practice questions for CBSE Exams as per the latest pattern and marking scheme issued by CBSE for the academic session 2019-20.

**Series 6**

**The p-Block Elements**

**Organic Chemistry – Some  
Basic Principles and Techniques**

Time Allowed : 3 hours  
Maximum Marks : 70

### GENERAL INSTRUCTIONS

- All questions are compulsory.
- Section A: Q.no. 1 to 20 are very short answer questions (objective type) and carry 1 mark each.
- Section B: Q.no. 21 to 27 are short answer questions and carry 2 marks each.
- Section C: Q.no. 28 to 34 are long answer questions and carry 3 marks each.
- Section D: Q.no. 35 to 37 are also long answer questions and carry 5 marks each.
- There is no overall choice. However an internal choice has been provided in two questions of two marks, two questions of three marks and all the three questions of five marks weightage. You have to attempt only one of the choices in such questions.
- Use log tables if necessary, use of calculators is not allowed.

### SECTION - A

Read the given passage and answer the questions 1 to 5 that follow :

Boron forms a number of hydrides having the general formula  $B_nH_{n+4}$  and  $B_nH_{n+6}$ . These hydrides are called boranes. The simplest hydride of boron is diborane,  $B_2H_6$ . Boranes contain special types of bonds known as multicentre bonds. Boranes have high heat of combustion.

- How many 2-centered 2-electron bonds and 3-centered-2-electron bonds are there in the structure of diborane?
- Complete the following reaction :  
 $B_2H_6 + NH_3 \xrightarrow{\text{Heat}}$  (NCT 2014)
- Draw the structure of diborane.
- What happens when diborane undergoes reaction with carbon monoxide ?
- Give a laboratory method to prepare diborane.

Question 6 to 10 are one word answer :

- Which hybrid orbital is used by underlined carbon in the following molecule?  
 $CH_3-CH=\underline{C}H_2$
- Name the metal which is commonly used as a reducing agent in metallurgical operations.
- Write the IUPAC name of  
 $CH_3-CH_2-CH_2-CH_2-COOH$  (KVS 2017)
- Name the gas which is evolved in the Kjeldahl's method used for the estimation of nitrogen gas.
- Which is the basic building unit of all silicates?


Questions 11 to 15 are multiple choice questions :

- Which of the following statements is not correct regarding graphite and inorganic graphite?  
(a) Both are slippery in nature.  
(b) Both conduct electricity but graphite is less conducting in nature as compared to inorganic graphite



- (c) Graphite is not having any charge separation like inorganic graphite.  
 (d) All atoms in graphite as well as in inorganic graphite are  $sp^2$  hybridized.

12. The descending order of stability of the following carbonium ions is



- (a)  $\text{IV} > \text{II} > \text{I} > \text{III}$  (b)  $\text{II} > \text{IV} > \text{III} > \text{I}$   
 (c)  $\text{II} > \text{IV} > \text{I} > \text{III}$  (d)  $\text{IV} > \text{II} > \text{III} > \text{I}$

13. Oxalic acid reacts with concentrated  $\text{H}_2\text{SO}_4$  to give a mixture of two gases. When this mixture is passed through caustic potash, one of the gases is absorbed. What is the product formed by the absorbed gas with caustic potash?

- (a)  $\text{K}_2\text{SO}_4$  (b)  $\text{KHCO}_3$   
 (c)  $\text{K}_2\text{CO}_3$  (d)  $\text{KOH}$

14. The reaction,  
 $\text{C}_6\text{H}_6 + \text{Br}_2 \xrightarrow{\text{FeBr}_3} \text{C}_6\text{H}_5\text{Br} + \text{HBr}$  is

- (a) an electrophilic addition reaction  
 (b) a nucleophilic substitution reaction  
 (c) an electrophilic substitution reaction  
 (d) a nucleophilic addition reaction.

15. The substance used as an adsorbent in the column chromatography is


- (a)  $\text{Na}_2\text{O}$  (b)  $\text{Na}_2\text{SO}_4$  (c)  $\text{Al}_2\text{O}_3$  (d) alum.

#### Questions 16 to 20 :

- (a) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.  
 (b) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.  
 (c) Assertion is correct, but reason is wrong statement.  
 (d) Assertion is wrong, but reason is correct statement.

16. **Assertion :** In the case of polyfunctional compounds, the choice of principal functional group is made on the basis of order of preference.

**Reason :** The order of decreasing priority for some functional groups is



17. **Assertion :** Zeolites are the three-dimensional network of aluminosilicates.

**Reason :** Negative charge on zeolite structure is neutralised by positively charged  $\text{Al}^{3+}$  ions.

18. **Assertion :** Paper chromatography is a type of partition chromatography.

**Reason :** Moving phase is liquid and stationary phase is solid.

19. **Assertion :** Sn in +2 oxidation state is a reducing agent while Pb in +4 state is an oxidising agent.

**Reason :** Inert pair effect is due to participation of s-electrons in bond formation.

20. **Assertion :** The order of stability of carbocations is  $3^\circ > 2^\circ > 1^\circ$ .

**Reason :** Carbon atom in carbocation is in  $sp^3$  state of hybridisation.


#### SECTION - B

21. Suggest a reason as to why CO is poisonous.

(NCERT, KVS 2013, 2014) (HOTS)

OR

Complete the reactions :



22. Explain why  $\text{CH}_2=\text{CH}^-$  is a better nucleophile than  $\text{CH}\equiv\text{C}^-$ .


23. What are silicones? How are they prepared?

24. How many metamers of 3-pentanone are possible? Write their structures and IUPAC names.


OR

- (a) In which C—C bond of  $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ , the inductive effect is expected to be the least?

- (b) Explain why the following two structures, I and II cannot be the major contributors to the real structure of  $\text{CH}_3\text{COOCH}_3$ .



25. Describe what happens when boric acid is heated.  
 26. Explain the principle of paper chromatography.  
 27. Which of the following ions is more stable? Give reason.



#### SECTION - C

28. Give reasons :

- (a) Conc.  $\text{HNO}_3$  can be stored in aluminium container.  
 (b) A mixture of dilute  $\text{NaOH}$  and aluminium pieces is used to open drain.  
 (c) Aluminium alloys are used to make aircraft body.







OR

Give the reaction for the following :

- (i) Silicon is heated with methyl chloride at high temperature in the presence of copper.
- (ii) Silicon dioxide is treated with hydrogen fluoride.
- (iii) Hydrated alumina is treated with aqueous NaOH solution.

29. Arrange the following in order of property mentioned against each.



30. 0.35 g of an organic substance was Kjeldahlised and the ammonia obtained was passed into 100 mL of  $N/5 \text{ H}_2\text{SO}_4$ . The excess acid required 154 mL of  $N/10 \text{ NaOH}$  for neutralisation. Calculate the percentage of nitrogen in the compound.

OR

In a Dumas nitrogen estimation method, 0.30 g of an organic compound gave  $50 \text{ cm}^3$  of  $\text{N}_2$  collected at 300 K and 715 mmHg pressure. Calculate the percentage composition of nitrogen in the compound. (Vapour pressure of water at 300 K is 15 mmHg.)

(NCERT, NCT 2008, KVS 2017)

31. Starting from  $\text{SiCl}_4$ , prepare the following :

- (i) silicon
- (ii) silica gel
- (iii)  $\text{Na}_2\text{SiO}_3$

32. Identify the following types of organic reactions :

- (a)  $\text{CH}_3\text{CH}_2\text{Br} + \text{HS}^- \longrightarrow \text{CH}_3\text{CH}_2\text{SH} + \text{Br}^-$
- (b)  $(\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{HCl} \longrightarrow (\text{CH}_3)_2\text{ClC}-\text{CH}_3$
- (c)  $\text{CH}_3\text{CH}_2\text{Br} + \text{HO}^- \longrightarrow \text{CH}_2=\text{CH}_2 + \text{H}_2\text{O} + \text{Br}^-$

33. Account for the following :

- (a)  $\text{BCl}_3$  exist as a monomer while  $\text{AlCl}_3$  exist as a dimer.
- (b)  $\text{AlCl}_3$  exists but  $\text{TlCl}_3$  does not.

34. (i) Define hyperconjugation effect.

- (ii) Identify the nucleophiles from the following :  $\text{R}_3\text{N}$ ,  $>\text{C}=\text{O}$ ,  $\text{NC}^-$

- (iii) Why is hyperconjugation called no-bond resonance?

#### SECTION - D

35. (a) Draw the resonance structures of the following compounds :

- (i)  $\text{CH}_2=\text{CH}-\ddot{\text{Cl}}:$
- (ii)  $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$
- (iii)  $\text{CH}_2=\text{CH}-\underset{\text{H}}{\underset{|}{\text{C}}}=\text{O}$

(b) Identify the most stable species in the following set of ions by giving reasons :

- (i)  $\overset{+}{\text{C}}\text{H}_3$ ,  $\overset{+}{\text{C}}\text{H}_2\text{Br}$ ,  $\overset{+}{\text{C}}\text{HBr}_2$ ,  $\overset{+}{\text{C}}\text{Br}_3$
- (ii)  $\bar{\text{C}}\text{H}_3$ ,  $\bar{\text{C}}\text{H}_2\text{Cl}$ ,  $\bar{\text{C}}\text{HCl}_2$ ,  $\bar{\text{C}}\text{Cl}_3$

OR

(a) Write the name of the isomerism shown by the following pairs :

- (i) buta-1,3-diene and but-1-yne
- (ii) ethoxybutane and propoxypropane

(b) Why  $3^\circ$  carbocations are more stable than  $1^\circ$  carbocations?

(c) Compare inductive and electromeric effects.

(d) Why  $\text{CCl}_3\text{COOH}$  is a stronger acid than  $(\text{CH}_3)_3\text{CCOOH}$ ? (KVS 2016)

36. Compound (X) on reduction with  $\text{LiAlH}_4$  gives a hydride (Y) containing 21.72% hydrogen along with other products. The compound (Y) reacts with air explosively resulting in boron trioxide. Identify (X) and (Y). Give balanced reactions involved in the formation of (Y) and its reaction with air.

OR

Describe the general trends for the following properties of groups 13 elements.

- (i) Atomic size
- (ii) Ionisation enthalpy
- (iii) Metallic character
- (iv) Oxidation states
- (v) Nature of halides.

37. (a) What is the difference between distillation, distillation under reduced pressure and steam distillation?



(b) Why is nitric acid added to sodium extract before adding silver nitrate for testing halogens?

OR

(a) Draw structures of all isomeric ethers corresponding to the molecular formula  $\text{C}_5\text{H}_{12}\text{O}$ .


(b) Give IUPAC names for the following compounds :







### SOLUTIONS


1. The four terminal B—H bonds are regular two centred-two electron bonds while the two bridge (B—H—B) bonds are different and can be described as three centred-two electron bonds.



3.



5. A laboratory method for the preparation of diborane involves the oxidation of sodium borohydride with iodine.



6.  $sp^2$

7. Aluminium

8. Pentanoic acid

9. Ammonia gas

10.  $\text{SiO}_4^{4-}$




11. (b)

12. (c) : Stability of carbocations is enhanced by

(i) conjugation

(ii) and presence of groups having +I effect.

So, the correct order is



14. (c)

15. (c)

16. (a)

17. (c) : Zeolites are the three-dimensional aluminosilicates in which few Si atoms are replaced by Al atoms. The acquired negative charge on zeolite structure is balanced by cations such as  $\text{Na}^+$ ,  $\text{K}^+$  or  $\text{Ca}^{2+}$ , etc.


18. (c)

19. (c) : In heavier elements, the tendency to show +2 oxidation state increases in the sequence  $\text{Ge} < \text{Sn} < \text{Pb}$ . It is due to the inability of  $ns^2$  electrons of valence shell to participate in bonding (inert pair effect).

20. (c)

21. The poisonous nature of carbon monoxide (CO) is due to its ability to form a complex with haemoglobin which is about 300 times more stable than the oxygen-haemoglobin complex. This (CO-haemoglobin complex) prevents haemoglobin (in the red blood corpuscles) from carrying oxygen in the body and ultimately resulting in death.


OR



22. In  $\text{H}_2\text{C}=\text{CH}^-$ , the carbon atom carrying the -ve charge is  $sp^2$ -hybridized while in  $\text{CH}\equiv\text{C}^-$  the carbon atom carrying the -ve charge is  $sp$ -hybridized. Since a  $sp^2$ -hybridized carbon is less electronegative than a  $sp$ -hybridized carbon, therefore,  $\text{H}_2\text{C}=\text{CH}^-$  is a better nucleophile than  $\text{HC}\equiv\text{C}^-$ .

23. They are organosilicon polymers having Si - O - Si linkages, the main unit being  $\text{R}_2\text{SiO}$ , (where R = alkyl or aryl groups). They are called silicones as their empirical formula ( $\text{R}_2\text{SiO}$ ) resembles that of ketones ( $\text{R}_2\text{CO}$ ). Silicones are prepared by hydrolysis of alkyl or aryl substituted chlorosilanes and their subsequent polymerisation.

24. Two metamers (I and II) of 3-pentanone are possible :



OR

(a) Magnitude of inductive effect diminishes as the number of intervening bonds increases. Hence, the effect is least in the bond between carbon-2 and carbon-3.

(b) The two structures are less important contributors as they involve charge separation. Additionally, structure (I) contains a carbon atom with an incomplete octet.

25. Boric acid on heating above 370 K leads to partial removal of water to yield metaboric acid,  $\text{HBO}_2$ . Further








**26.** The underlying principle of paper chromatography is that of partition chromatography which is based on continuous differential partitioning of components of mixture between stationary and mobile phases. In paper chromatography, the paper used has water trapped in it which acts as the stationary phase while a suitable solvent or a mixture of solvents is used as a mobile phase.

(A)  $\longleftrightarrow$  (B)

**28. (a)** Conc.  $\text{HNO}_3$  can be stored in aluminium container because conc.  $\text{HNO}_3$  makes aluminium passive due to the formation of a protective layer of  $\text{Al}_2\text{O}_3$  on its surface.

(c) Aluminium alloys are used for making aircraft body because these are light in weight, corrosion resistant, strong and highly ductile.

(i)  $\text{CH}_3\text{Cl} + \text{Si} \xrightarrow[570 \text{ K}]{\text{Cu powder}} \text{CH}_3\text{SiCl}_3 + (\text{CH}_3)_2\text{SiCl}_2 + (\text{CH}_3)_3\text{SiCl} + (\text{CH}_3)_4\text{Si}$   
Methyl chloride






$$\frac{N}{5} \times V_{\text{acid}} = \frac{N}{10} \times 154 \text{ mL}$$

$$\frac{1.4 \times \text{Normality of acid} \times \text{Volume of acid used for neutralising NH}_3}{\text{Mass of the compound taken}}$$

$$\text{Volume of nitrogen at STP} = \frac{273 \times 700 \times 50}{300 \times 760} = 41.9 \text{ mL}$$

$$41.9 \text{ mL of nitrogen will weigh} = \frac{28 \times 41.9}{22400} \text{ g}$$

$$\% \text{ of nitrogen} = \frac{28 \times 41.9 \times 100}{22400 \times 0.3} = 17.46\%$$



**(b)** In the given reaction, we see that the  $\text{H}^+$  and  $\text{Cl}^-$  reacted with the alkene and thus, produced a haloalkane. However, no atom has been displaced. Thus, the reaction is an electrophilic addition reaction.

**30** ● ● **CHEMISTRY TODAY** | DECEMBER '19



**33. (a)** Due to small size of boron, there is a dative  $p\pi - p\pi$  bonding takes place between the lone pair of Cl atoms and vacant  $p$ -atomic orbital of B. Hence,  $\text{BCl}_3$  is a monomer.




Due to large size, Al cannot participate in dative  $p\pi - p\pi$  bonding. Therefore, in order to complete the octet,  $\text{AlCl}_3$  dimerizes.

**(b)** Because as we move down the group, the tendency of  $ns^2$  electron to remain inert increases (inert pair effect). Therefore, only one electron from the  $p$ -atomic orbital participates in the bond formation and hence +1 oxidation state becomes more and more stable as compared to +3 oxidation state. Therefore, Tl forms only  $\text{TlCl}$ .

**34. (i)** When an alkyl group is attached to an unsaturated system such as a double bond or a benzene ring, the order of inductive effect is actually reversed. This effect is called hyperconjugation effect or Baker-Nathan effect.

**(ii)**  $\text{NC}^-$

**(iii)** Hyperconjugation is called no-bond resonance because the hydrogen bonded to carbon loses its bond from carbon during resonance.




**(b) (i)**  $\overset{+}{\text{CH}_3}$  is most stable because the replacement of hydrogen by bromine increases positive charge on carbon atom and destabilises the species.


**(ii)**  $\ddot{\text{CCl}_3}$  is most stable because electronegativity of chlorine is more than hydrogen. Negative charge on carbon is reduced which results in more stability.

OR

**(a) (i) Functional isomerism :**




**(ii) Metamerism :**




**(b)** Hyperconjugation interaction in  $(\text{CH}_3)_3\overset{+}{\text{C}}$  is greater than in  $\text{CH}_3\overset{+}{\text{CH}_2}$  as  $(\text{CH}_3)_3\overset{+}{\text{C}}$  has nine C—H bonds.



**(c)** Inductive effect is a permanent effect caused due to more or less electronegative atom/group attached to carbon in an organic compound.




Electromeric is a temporary electron displacement caused in carbon chain due to presence of an external electrophile or nucleophile.



**(d)**  $\text{CCl}_3\text{COOH}$  is a stronger acid than  $(\text{CH}_3)_3\text{CCOOH}$  due to presence of three electron withdrawing groups ( $I$ -effect). This reduces electron density of —OH bond and hence, make it more acidic.



From this we can guess 'Y' must be  $\text{B}_2\text{H}_6$ .




% of hydrogen in  $\text{B}_2\text{H}_6 = \frac{6}{27.62} \times 100 = 21.72\%$

[Mol. wt. of  $\text{B}_2\text{H}_6 = 21.62 + 6 = 27.62$ ]

Thus, percentage of hydrogen is 21.72 in  $\text{B}_2\text{H}_6$ .

From this we can guess that X is boron trihalide.



OR

**(i) Atomic size :** Atomic radii of group-13 elements increase down the group with the exception that atomic radius of Ga is less than that of Al due to the presence of  $d$ -electrons which offer poor screening effect for the outer electrons from the increased nuclear charge in Ga.

**(ii) Ionisation enthalpy :** For group-13 elements, the trend of ionisation enthalpy is  $\text{B} > \text{Al} < \text{Ga} > \text{In} < \text{Tl}$ . This is due to increase in size and low screening effect of  $d$ - and  $f$ -electrons.

**(iii) Metallic character :** Metallic character increases from boron to aluminium then decreases down the group.

**(iv) Oxidation states :** For group-13 elements, both +1 and +3 oxidation states are observed. The +1 oxidation state becomes more stable as we move down the group due to inert pair effect. Boron does not show +3 oxidation state.

**(v) Nature of halides :** Group-13 elements form trihalides (except  $\text{TlI}_3$ ). Due to electron deficient nature,  $\text{BCl}_3$  accepts electrons and forms adducts.  $\text{AlCl}_3$  achieves stability by forming a dimer.



37. (a) Differences between distillation, distillation under reduced pressure and steam distillation :

Criterion for difference	Distillation	Distillation under reduced pressure	Steam distillation
Principle	It is used to separate – volatile liquids from non-volatile impurities. – liquids having sufficient difference in their boiling points.	It is used to purify liquids – having very high boiling points. – which decompose at or below their boiling point	This method is used to separate substances which are steam volatile, immiscible with water, possess a vapour pressure of 10–15 mm Hg and contain non-volatile impurities
Used for	Separation of volatile liquids from non-volatile impurities.	Separation of liquids having very high boiling points or those, which decompose at or below their boiling points.	For a substances which are immiscible in water but are steam volatile.
Example	A mixture of chloroform (b.pt. 334 K) and aniline (b.pt. 457 K).	Separation of glycerol from spent – lye in soap industry.	Separation of aniline from aniline-water mixture.

(b) If N or S is present in the compound, the sodium fusion extract is first boiled with concentrated nitric acid to decompose cyanide or sulphide of sodium during Lassaigne's test. These ions would otherwise interfere with silver nitrate test for halogens.

OR

(a) The possible isomeric ethers corresponding to  $C_5H_{12}O$  are,

(i)	$C_2H_5-O-C_3H_7$	$  \begin{array}{ccccccc}  & H & H & & H & H & H \\  &   &   & &   &   &   \\  H & -C & -C & -O & -C & -C & -C-H \\  &   &   & &   &   &   \\  & H & H & & H & H & H  \end{array}  $
(ii)	$CH_3-O-C_4H_9$	$  \begin{array}{ccccccc}  & H & & H & H & H & H \\  &   & &   &   &   &   \\  H & -C & -O & -C & -C & -C & -C-H \\  &   & &   &   &   &   \\  & H & & H & H & H & H  \end{array}  $
(iii)	$CH_3CH_2-O-CH(CH_3)_2$	$  \begin{array}{ccccccc}  & & & H & & & \\  & & &   & & & \\  & H & H & H-C-H & H & & \\  &   &   &   &   & & \\  H & -C & -C & -O & -C & -C & -H \\  &   &   & &   & &   \\  & H & H & & H & & H  \end{array}  $
(iv)	$  \begin{array}{c}  CH_3 \\    \\  CH_3-O-C-CH_3 \\    \\  CH_3  \end{array}  $	$  \begin{array}{ccccccc}  & & & H & & & \\  & & &   & & & \\  & H & H & H-C-H & H & & \\  &   &   &   &   & & \\  H & -C & -O & -C & -C & -H \\  &   & &   &   & & \\  & H & & H-C-H & H & & \\  & & &   & & & \\  & & & H & & &   \end{array}  $


(b) (i) Cyclohexane-1, 2-diol

(ii) 6-Chloro-2-methylhexan-3-one





# MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 120

## Some Basic Concepts of Chemistry | Structure of Atom

Time Taken : 60 Min.

NEET / AIIMS

Only One Option Correct Type

- A solution of KCl has a density of  $1.69 \text{ g mL}^{-1}$  and is 67% by weight. Find the density ( $\text{g mL}^{-1}$ ) of the solution if it is diluted so that the percentage by weight of KCl in the diluted solution is 30%.  
(a) 1.224 (b) 2.22  
(c) 5.68 (d) 4.52
- An electron in H atom jumps from the third energy level to the first energy level. The change in the potential energy of the electron is  
(a) 12.09 eV (b) 6.04 eV  
(c) 24.18 eV (d) 9.24 eV
- When 3.2 g S is vaporized at  $450^\circ\text{C}$  and 723 mm pressure, the vapours occupy a volume of 780 mL. If the molecular formula of S vapour under these conditions is  $\text{S}_x$ , then the value of x is  
(a) 6 (b) 4  
(c) 2 (d) 8
- An electron beam can undergo diffraction by crystals. Through what potential should a beam of electrons be accelerated so that its wavelength becomes equal to  $1.54 \text{ \AA}$ ?  
(a) 5.6 V (b) 24.5 V  
(c) 42.4 V (d) 63.4 V
- An aqueous solution of 6.3 g oxalic acid dihydrate is made up to 250 mL. The volume of 0.1 N NaOH required to completely neutralise 10 mL of this solution is  
(a) 40 mL (b) 20 mL  
(c) 10 mL (d) 4 mL.
- The kinetic energy of an electron in the second Bohr orbit of a hydrogen atom is  
(a)  $\frac{h^2}{4\pi^2 m a_0^2}$  (b)  $\frac{h^2}{16\pi^2 m a_0^2}$   
(c)  $\frac{h^2}{32\pi^2 m a_0^2}$  (d)  $\frac{h^2}{64\pi^2 m a_0^2}$
- When 10 mL of ethyl alcohol (density =  $0.7893 \text{ g mL}^{-1}$ ) is mixed with 20 mL of water (density  $0.9971 \text{ g mL}^{-1}$ ) at  $25^\circ\text{C}$ , the final solution has a density of  $0.9571 \text{ g mL}^{-1}$ . The percentage change in total volume on mixing is  
(a) 3.1% (b) 2.4%  
(c) 1% (d) none of these.
- The given electronic configuration deviates from  

$$\begin{array}{|c|c|c|c|} \hline ns & np & & \\ \hline \uparrow & \uparrow\uparrow & \uparrow & \square \\ \hline \end{array}$$
  
(I) Hund's rule  
(II) Aufbau principle  
(III) Pauli's principle  
(a) All of the above (b) Only I, II  
(c) Only I, III (d) Only II, III
- Sulphuric acid reacts with sodium hydroxide as follows :  

$$\text{H}_2\text{SO}_4 + 2\text{NaOH} \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$$
  
When 1 L of 0.1 M sulphuric acid solution is allowed to react with 1 L of 0.1 M sodium hydroxide solution then the amount of sodium sulphate formed and its molarity in the solution obtained is



- (a) 7.10 g and  $0.1 \text{ mol L}^{-1}$   
 (b) 7.10 g and  $0.025 \text{ mol L}^{-1}$   
 (c) 5.40 g and  $0.1 \text{ mol L}^{-1}$   
 (d) 3.55 g and  $0.025 \text{ mol L}^{-1}$ .

10. When electronic transition occurs from higher energy state to lower energy state with energy difference equal to  $\Delta E$  electron volts, the wavelength of the line emitted is approximately equal to ( $h = 6.62 \times 10^{-34} \text{ Js}$ )

- (a)  $\frac{12397}{\Delta E} \times 10^{-10} \text{ m}$  (b)  $\frac{12397}{\Delta E} \times 10^{10} \text{ m}$   
 (c)  $\frac{12397}{\Delta E} \times 10^{-10} \text{ cm}$  (d)  $\frac{12397}{\Delta E} \times 10^{10} \text{ cm}$

11. Two oxides of a certain metal were separately heated in a current of hydrogen until constant weights were obtained. The water produced in each case was carefully collected and weighed. 2 g of each oxide gave, respectively 0.2517 g and 0.4526 g of water. This observation illustrates  
 (a) law of conservation of mass  
 (b) law of constant proportions  
 (c) law of multiple proportions  
 (d) law of reciprocal proportions.

12. The decreasing order of energy for an electrons represented by the following sets of quantum number is

1.  $n = 4, l = 0, m = 0, s = \pm 1/2$   
 2.  $n = 3, l = 1, m = 1, s = -1/2$   
 3.  $n = 3, l = 2, m = 0, s = +1/2$   
 4.  $n = 3, l = 0, m = 0, s = -1/2$   
 (a)  $1 > 2 > 3 > 4$  (b)  $2 > 1 > 3 > 4$   
 (c)  $3 > 1 > 2 > 4$  (d)  $4 > 3 > 2 > 1$

#### Assertion & Reason Type

**Directions :** In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.  
 (b) If both assertion and reason are true but reason is not the correct explanation of assertion.  
 (c) If assertion is true but reason is false.  
 (d) If both assertion and reason are false.

13. **Assertion :** If 100 mL of 0.2 N HCl is mixed with 100 mL of 0.3 N HCl, the normality of final solution will be 0.25 N.

**Reason :** If two solutions of the same solute are mixed, the normalities can be added.

14. **Assertion :** Orbitals with the same value of  $l$  and  $m_l$  have the same angular wave function.

**Reason :** Angular wave function doesn't depend upon the  $n$ .

15. **Assertion :** For a solution, 92 g of ethanol and 144 g of water is mixed, mole fraction of ethanol and water in the solution is 0.2 and 0.8 respectively.

**Reason :** The sum of the mole fractions of all the components in a solution is unity.

#### JEE MAIN / ADVANCED

#### Only One Option Correct Type

16. Photoelectric emission is observed from a surface for frequencies  $\nu_1$  and  $\nu_2$  of the incident radiation ( $\nu_1 > \nu_2$ ). If the maximum kinetic energies of the photo-electrons in the two cases are in the ratio  $1 : k$  then the threshold frequency  $\nu_0$  is given by

- (a)  $\frac{\nu_2 - \nu_1}{k - 1}$  (b)  $\frac{k\nu_1 - \nu_2}{k - 1}$   
 (c)  $\frac{k\nu_2 - \nu_1}{k - 1}$  (d)  $\frac{\nu_2 - \nu_1}{k}$

17. 1.2 g mixture of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  was dissolved in water to form  $100 \text{ cm}^3$  of a solution.  $20 \text{ cm}^3$  of this solution required  $40 \text{ cm}^3$  of 0.1 N HCl for neutralisation. What will be the weight of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$  in the mixture ?

- (a)  $\text{Na}_2\text{CO}_3 = 0.596 \text{ g}$ ,  $\text{K}_2\text{CO}_3 = 0.604 \text{ g}$   
 (b)  $\text{Na}_2\text{CO}_3 = 1.56 \text{ g}$ ,  $\text{K}_2\text{CO}_3 = 2.46 \text{ g}$   
 (c)  $\text{Na}_2\text{CO}_3 = 3.52 \text{ g}$ ,  $\text{K}_2\text{CO}_3 = 5.6 \text{ g}$   
 (d)  $\text{Na}_2\text{CO}_3 = 4.56 \text{ g}$ ,  $\text{K}_2\text{CO}_3 = 0.24 \text{ g}$

18. The atomic spectrum of hydrogen is found to contain a series of lines at wavelengths (in nm) 656.46, 486.27, 434.17 and 410.29. What will be the wavelength of the next line in the series?

- (a) 2.56 nm (b) 6.52 nm  
 (c) 1.592 nm (d) 397.1 nm

19. The mass of one litre sample of ozonised oxygen at N.T.P. was found to be 1.5 g. When 100 mL of this mixture at N.T.P. were treated with turpentine oil, the volume was reduced to 90 mL. The molecular mass of ozone is

- (a) 98 (b) 48 (c) 56 (d) 68

#### More than One Options Correct Type

20. Choose the correct statement(s) regarding the photoelectric effect.

- (a) No electrons are ejected, regardless of the intensity of the radiation, unless the frequency



exceeds a threshold value characteristic of the metal.

- (b) The kinetic energy of the ejected electrons varies linearly with frequency of the incident radiation and its intensity.
- (c) Even at low intensities, electrons are ejected immediately if the frequency is above the threshold value.
- (d) An intense and weak beam of monochromatic radiations differ in having number of photons and not in the energy of photons.

21. Which of the following statements are correct?

- (a) A sample of  $\text{CaCO}_3$  contains  $\text{Ca} = 40\%$ ,  $\text{C} = 12\%$ , and  $\text{O} = 48\%$ . If the law of constant composition is true, then the mass of  $\text{Ca}$  in 10 g of  $\text{CaCO}_3$  from another source is 4.0 g.
- (b) 12 g of carbon is heated in vacuum and there is no change in the mass, is the best example of the law of conservation of mass.
- (c) Air is heated at constant pressure and there is no change in mass but the volume increases, is the best example of the law of conservation of mass.

- (d)  $\text{SO}_2$  gas was prepared by (i) heating  $\text{Cu}$  with conc.  $\text{H}_2\text{SO}_4$ , (ii) burning sulphur in oxygen, (iii) reacting sodium sulphite ( $\text{Na}_2\text{SO}_3$ ) with dilute  $\text{H}_2\text{SO}_4$ . It was observed that in each case,  $\text{S}$  and  $\text{O}$  combine in the ratio of 1 : 1. This data illustrates the law of constant composition.

22. Which of the following statements are correct?

- (a) Energy of an orbital is determined by principal quantum number.
- (b) For all values of  $n$ , the  $p$ -orbital have the same shape, but overall size increases as  $n$  increases, for a given atom.
- (c) A  $2p_x$  atomic orbital consists of two lobes of electron density.
- (d) Magnetic quantum number  $= \sqrt{n(n+2)}$  B.M.

23. 1.24 g of  $\text{P}_4$  reacts with 8.0 g of  $\text{O}_2$ .

The following reactions occur :

- (i)  $\text{P}_4 + 5\text{O}_2 \longrightarrow \text{P}_4\text{O}_{10}$
- (ii)  $\text{P}_4 + 3\text{O}_2 \longrightarrow \text{P}_4\text{O}_6$

Which of the following statements are wrong?

- (a)  $\text{P}_4$  is the limiting quantity.
- (b)  $\text{O}_2$  is the limiting quantity.
- (c) Mass of  $\text{P}_4\text{O}_{10}$  obtained is 2.2 g.
- (d) Mass of  $\text{P}_4\text{O}_6$  obtained is 2.84 g.

## Scientist who Made Us Proud

Subramania Ranganathan (1934–2016), popularly known as Ranga, was an Indian bioorganic chemist and a professor and head of the department of chemistry at the Indian Institute of Technology, Kanpur. He was known for his studies on synthetic and mechanistic organic chemistry and was an elected fellow Indian National Science Academy, National Academy of Sciences, India and the Indian Academy of Sciences.



**Subramania Ranganathan**  
(2 Feb 1934 - 8 Jan 2016)

### Biography

Ranganathan, born on 2 February 1934 in the south Indian state of Tamil Nadu, graduated in chemistry from Madras University and continued there to complete his master's degree. In the US, he enrolled at Ohio State University at Harold Shechter's laboratory and secured a PhD in 1962. He moved to the laboratory of Robert Burns Woodward, the 1965 Nobel laureate, at Harvard University for his post-doctoral studies and in 1964, he shifted to Woodward Research Institute, Basel to complete the studies in 1964. In 1966, he joined IIT Kanpur where he spent his entire official academic career, holding positions of a professor, head of the department and dean, before superannuating in

1994. Post-retirement, he served as an INSA senior scientist, first at National Institute for Interdisciplinary Science and Technology and later at the Indian Institute of Chemical Technology (IICT).

Ranganathan was holding the position of an honorary position at IICT when he died on 8 January 2016, at the age of 81.

During his post-doctoral days, Ranganathan worked closely with Woodward and was known to have assisted the latter in his work on Woodward–Hoffmann rules.

### Awards and Honors

- Ranganathan received the Basudev Banerjee Medal in 1975 and the Council of Scientific and Industrial Research awarded him the Shanti Swarup Bhatnagar Prize in 1977.
- He received R. C. Mehrotra Endowment Gold Medal in 2000 and the Silver Medal of the Chemical Research Society of India in 2001.
- CRSI honored him again in 2006 with the Lifetime Achievement Award.
- In 2014, he was awarded the Best Teacher Award by the Indian National Science Academy.
- The Indian Academy of Sciences elected him as a fellow in 1975 and he became an elected fellow of the Indian National Science Academy and the National Academy of Sciences, India in 1981 and 1991 respectively.



### Numerical Value Type

24. A mixture of formic acid and oxalic acid is heated with conc.  $\text{H}_2\text{SO}_4$ . The gas produced is collected and on its treatment with KOH solution, the volume of the gas decreased by  $1/6^{\text{th}}$ . If the molar ratio of the two acids in the original mixture is  $x : 1$ , then find the value of  $x$ .
25. If the position of the electron is measured within an accuracy of  $\pm 0.002 \text{ nm}$ , the uncertainty in the momentum of the electron is  $x \times 10^{-23} \text{ kg ms}^{-1}$ , then the value of  $x$  is
26. A 100% pure sample of a divalent metal carbonate weighing 2 g on complete thermal decomposition releases 448 cc of carbon dioxide at STP. The equivalent mass of the metal is

### Matrix Match Type

Answer the following questions (27 and 28) by appropriately matching the columns based on the information given in the passage :

The set of an electron is described by a set of four quantum numbers.

- Principal quantum number ( $n$ ) gives the size of the shell and the relative average distance of an electron from the nucleus and thereby the energy of the electron.
- Azimuthal quantum number or subsidiary quantum number ( $l$ ) gives the subshell and shape of the orbital for an electron. Each orbital of a given shell is degenerate.  $l = 0, 1, 2, 3, 4, \dots$  indicates the subshell  $s, p, d, f, g, \dots$  respectively. A subshell is only possible when  $l < n$  and total subshells in a shell is equal to numerical value of  $n$ .
- Magnetic quantum number ( $m$ ) determines the preferred orientations of orbitals in space.  $m = -l, \dots, 0, \dots, +l$ .
- Spin quantum number ( $s$ ) represents the spin of an electron. In an orbital there are maximum of two electrons such that their spins are  $+1/2$  and  $-1/2$  and the sum of the spins is zero.

Column-I		Column-II	
P.	$n = 2, l = 1, m = \pm 1$	I.	$3d_{x^2-y^2}$ or $3d_{xy}$
Q.	$n = 4, l = 2, m = 0$	II.	$2p_x$ or $2p_y$
R.	$n = 4, l = 0, m = 0$	III.	$4s$
S.	$n = 3, l = 2, m = \pm 2$	IV.	$4d_z^2$

27. Which of the following has the correct combination considering column I and column II ?  
 (a)  $P \rightarrow \text{II}$  (b)  $Q \rightarrow \text{III}$   
 (c)  $R \rightarrow \text{I}$  (d)  $S \rightarrow \text{IV}$
28. Which of the following has correct combination considering column I and column II ?  
 (a)  $P \rightarrow \text{IV}$  (b)  $Q \rightarrow \text{I}$  (c)  $R \rightarrow \text{III}$  (d)  $S \rightarrow \text{II}$

Answer the following questions (29 and 30) by appropriately matching the columns based on the information given in the passage :

In a balanced chemical reaction the reactant which is present in the lesser amount gets consumed after sometime and after that no further reaction takes place whatever be the amount of the other reactants present. Hence, the reactant which gets consumed, limits the amount of product formed and is, therefore, called the limiting reagent.

Column-I		Column-II	
P.	$2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$ 1.0 g 1.0 g ?	I.	0.56 g
Q.	$\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$ 1.0 g 1.0 g ?	II.	1.333 g
R.	$\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$ 1.0 g ?	III.	1.125 g
S.	$\text{C} + 2\text{H}_2 \rightarrow \text{CH}_4$ 1.0 g 1.0 g ?	IV.	1.214 g

29. Which of the following has correct combination considering column I and column II ?  
 (a)  $P \rightarrow \text{III}$  (b)  $Q \rightarrow \text{I}$  (c)  $R \rightarrow \text{II}$  (d)  $S \rightarrow \text{IV}$
30. Which of the following has correct combination considering column I and column II ?  
 (a)  $P \rightarrow \text{II}$  (b)  $Q \rightarrow \text{III}$   
 (c)  $R \rightarrow \text{I}$  (d)  $S \rightarrow \text{IV}$



Keys are published in this issue. Search now! 😊

## CHECK YOUR PERFORMANCE

No. of questions attempted .....

No. of questions correct .....

Marks scored in percentage .....

If your score is

> 80%

Your preparation is going good, keep it up to get high score.

60-80%

Need more practice, try hard to score more next time.

< 60%

Stress more on concepts and revise thoroughly.



# GET SET GO for JEE




*with exclusive and brain storming MCQs*

Practicing these MCQs help to strengthen your concepts and give you extra edge in your JEE preparation


## Only One Option Correct Type

- Which of the following is true for the species having  $3d^4$  configuration?  
 (a)  $\text{Cr}^{2+}$  is reducing in nature.  
 (b)  $\text{Mn}^{3+}$  is oxidising in nature.  
 (c) Both (a) and (b).  
 (d) None of these.

- The products (A) and (B) are respectively




- (a) and   
 (b) and   
 (c) and



- Which of the following statements is correct regarding enzymes?  
 (a) Enzymes are specific biological catalysts that can normally function at very high temperatures ( $T \sim 1000 \text{ K}$ ).  
 (b) Enzymes are normally heterogeneous catalysts that are very specific in action.  
 (c) Enzymes are specific biological catalysts that cannot be poisoned.  
 (d) Enzymes are specific biological catalysts that possess well defined active sites.
- An element X of atomic mass 25.0 exists as  $\text{X}_4$  in benzene to the extent of 100%. When 10.30 g of saturated solution of X in benzene is added to 20.0 g of benzene, the depression in freezing point of the resulting solution is 0.51 K. If  $K_f$  for benzene is  $5.1 \text{ K kg mol}^{-1}$ , the solubility of X in 100 g of benzene will be  
 (a) 3.0 g (b) 2.7 g  
 (c) 0.30 g (d) 0.27 g.




5. Compare the properties of two isomeric products X and Y formed in the following reaction :



Acid strength	H <sub>2</sub> O solubility	Volatility	Melting point
(a) Y > X	Y > X	X > Y	Y > X
(b) X > Y	X > Y	Y > X	X > Y
(c) Y > X	X > Y	Y > X	Y > X
(d) X > Y	Y > X	X > Y	Y > X

6. The arrangement of X<sup>-</sup> ions around A<sup>+</sup> ion in solid AX is given in the figure (not drawn to scale). If the radius of X<sup>-</sup> is 250 pm, the radius of A<sup>+</sup> is



- (a) 104 pm (b) 125 pm  
(c) 183 pm (d) 57 pm.

7. Identify the products.



- (a) Na<sub>2</sub>SO<sub>4</sub>, HCl (b) Na<sub>2</sub>S, HCl  
(c) NaClO<sub>3</sub>, S (d) Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, HCl

8. Which of the following complexes having tetrahedral geometry?

- (I) Ni(CO)<sub>4</sub>  
(II) K[AgF<sub>4</sub>]  
(III) Na<sub>2</sub>[Zn(CN)<sub>4</sub>]  
(IV) K<sub>2</sub>[PtCl<sub>4</sub>]  
(V) [RhCl(PPh<sub>3</sub>)<sub>3</sub>]

- (a) Only II, III and V (b) Only I, II and III  
(c) Only I, III and IV (d) Only I and III

9. Which one of the following is a chain growth polymer?

- (a) Polymethyl methacrylate  
(b) Nucleic acid  
(c) Polystyrene  
(d) Protein


10. Match column-I (metal) with column-II (procedure of extraction) and select the correct answer by choosing an appropriate option.

Column-I		Column-II	
A.	Pb	p.	Hydrometallurgical process
B.	Cu	q.	Carbon reduction method
C.	Fe	r.	Thermite process
D.	Cr	s.	Self-reduction method

- (a) A → q, s; B → r, s; C → q; D → r  
(b) A → q, s; B → p, q, r; C → p, q; D → p, r  
(c) A → p, s; B → p, q, r, s; C → p, q, s; D → p, q, r, s  
(d) A → p; B → q, r; C → r; D → s

### More Than One Options Correct Type

11. Which of the following statement are true regarding the product formed in the given reaction?




- (a) Major product is given by S<sub>N</sub>1 reaction.  
(b) Through E1 mechanism three alkenes are formed.  
(c) 3-Methylpentan-3-ol is also formed as one of the product.  
(d) Fractional distillation of elimination product will give two fractions.

12. Which of the following statements are not correct for lyophilic colloids?

- (a) These are prepared by special indirect methods.  
(b) These are not much hydrated.  
(c) These are also called intrinsic colloids.  
(d) Small quantity of electrolyte is sufficient to cause precipitation of these colloids.

### Numerical Value Type

13. If the standard reduction potential for Cu<sup>2+</sup>|Cu is +0.34 V, then the reduction potential at pH = 14 for the above couple is -xV. The value of x is (K<sub>sp</sub> of Cu(OH)<sub>2</sub> is 1.0 × 10<sup>-19</sup>)



The total number of halogen atoms present in the major product is


15. The rate of a first-order reaction is 0.04 mol litre<sup>-1</sup> s<sup>-1</sup> at 10 minutes and 0.03 mol litre<sup>-1</sup> s<sup>-1</sup> at 20 minutes



after initiation. The half-life (in min) of the reaction is

### SOLUTIONS

1. (c) :  $\text{Cr}^{2+}$  is reducing as its configuration changes from  $d^4$  to  $d^3$ , this  $d^3$  configuration can be written as  $t_{2g}^3$ , which is a more stable configuration. On the other hand, the change from  $\text{Mn}^{3+}$  to  $\text{Mn}^{2+}$  results in the half-filled ( $d^5$ ) configuration which has extra stability.



3. (d)
4. (a) : Suppose saturated solution of X in benzene contains  $w$  g of X (present as  $X_4$ ). Hence, amount of benzene present =  $(10.30 - w)$  g.

$\therefore$  Total amount of benzene present  
 $= 20 + (10.30 - w)\text{g} = (30.30 - w)\text{g}$

$$\Delta T_f = \frac{1000 \times K_f \times w_2}{w_1 \times M_2}$$

$$0.51 = \frac{1000 \times 5.1 \times w}{(30.30 - w) \times 100} \quad (M_2 \text{ of } X_4 = 25 \times 4 = 100)$$


$$\text{or } 51(30.30 - w) = 5100 w$$

$$\text{or } 30.30 - w = 100 w$$

$$\text{or } 101 w = 30.30 \text{ or } w = 0.3 \text{ g}$$

Thus, 10.30 g of saturated solution contains 0.3 g of X and 10 g of benzene.

$\therefore$  Solubility of X in 100 g of benzene = 3.0 g



Acid strength :  $X > Y$  (Carboxylate anion stabilized by H-bonding)


Solubility in  $\text{H}_2\text{O}$  :  $Y > X$  (Intermolecular H-bonding in Y)

Volatility :  $X > Y$  (Intramolecular H-bonding in X)

M.Pt. :  $Y > X$  (More symmetrical structure of Y)



### COMIC CAPSULE





6. (a): Cation  $A^+$  occupies octahedral void arrangement by anion  $X^-$ .

$$\frac{r_{A^+}}{r_{X^-}} = 0.414$$


$$\frac{r_{A^+}}{250} = 0.414 \quad [\because \text{Radius of } X^- (r_X^-) = 250 \text{ pm}]$$

$$r_{A^+} = 0.414 \times 250 = 103.5 \approx 104 \text{ pm}$$

7. (a):  $\text{Na}_2\text{SO}_3 + \text{Cl}_2 + \text{H}_2\text{O} \rightleftharpoons \text{Na}_2\text{SO}_4 + 2\text{HCl}$
8. (d): (I) It is tetrahedral ( $sp^3$ ) because there is no  $(n-1)d$  orbital vacant for  $dsp^2$  hybridisation.  
 (II) Ag with +3 oxidation state and  $4d^8$  configuration has square planar geometry.  
 (III) It is tetrahedral ( $sp^3$ ) because there is no  $(n-1)d$  orbital vacant for  $dsp^2$  hybridisation.  
 (IV) Pt with +2 oxidation state and  $5d^8$  configuration has square planar geometry.  
 (V) Rh with +1 oxidation state and  $4d^8$  configuration has square planar geometry.  
 (I) and (III) have tetrahedral geometry.

9. (c)


10. (a)



12. (a, b, d)

13. (0.22): At pH = 14,  $[\text{H}^+] = 1 \times 10^{-14} \text{ M}$ ;  
 $[\text{OH}^-] = 10^0 = 1 \text{ M}$   
 $\therefore [\text{H}^+][\text{OH}^-] = 1 \times 10^{-14}$

$\text{Cu}(\text{OH})_2$  ionises as follows:



$$\therefore K_{\text{sp}} \text{ of } \text{Cu}(\text{OH})_2 = [\text{Cu}^{2+}][\text{OH}^-]^2$$

$$\text{or } 1.0 \times 10^{-19} = [\text{Cu}^{2+}](1)^2$$

$$\text{or } [\text{Cu}^{2+}] = 1.0 \times 10^{-19} \text{ M}$$


The standard electrode potential of  $\text{Cu}^{2+}|\text{Cu}$  is represented as follows:  $\text{Cu}_{(\text{aq})}^{2+} + 2e^- \longrightarrow \text{Cu}_{(\text{s})}$

$$\text{Using Nernst equation, } E = E^\circ - \frac{0.059}{n} \log \frac{1}{[\text{Cu}^{2+}]}$$

$$= 0.34 - \frac{0.059}{2} \log \frac{1}{1 \times 10^{-19}} \quad [\because n = 2]$$

$$= 0.34 - \frac{0.059}{2} \log (10^{19}) = 0.34 - 0.0295 \times 19$$

$$= 0.34 - 0.56 = -0.22 \text{ V}$$



15. (24.15): Rate =  $kC$

$$\therefore \text{Rate, } r_1 = kC_1$$

[ $k$  is same]

$$\text{and rate } r_2 = kC_2$$

$$\therefore \frac{r_1}{r_2} = \frac{C_1}{C_2} \quad \therefore \frac{r_1 \text{ (at 10 min)}}{r_2 \text{ (at 20 min)}} = \frac{0.04}{0.03} = \frac{C_1}{C_2}$$

$$\text{Now } t = \frac{2.303}{k} \log \frac{C_1}{C_2} \quad [\text{For first order reaction}]$$

$$\text{or } k = \frac{2.303}{t} \log \frac{C_1}{C_2}$$

$$\text{When, } t = 10 \text{ min, } k = \frac{2.303}{10} \log \frac{0.04}{0.03}$$

$$\text{or } k = \frac{2.303}{10} \log \frac{4}{3} = 0.0287 \text{ min}^{-1}$$

$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{0.0287} = 24.15 \text{ min}$$





for

## NEET/JEE

Brush up your concepts to get high rank in NEET/JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

2020

Unit  
6

## Aldehydes, Ketones and Carboxylic Acids

## Aldehydes and Ketones

- **General formula :**  $C_nH_{2n}O$  having  $>C=O$  group. For ketones, value of  $n \geq 3$ .

– **Aldehydes :**  $\begin{array}{c} R \\ \diagdown \\ C=O \\ \diagup \\ H \end{array}$ ; where  $R = H$ , alkyl or aryl.


– **Ketones :**  $\begin{array}{c} R \\ \diagdown \\ C=O \\ \diagup \\ R \end{array}$ ; where  $R =$  alkyl or aryl.


- **Nomenclature :**


- The common names of most aldehydes are derived from the common names of the corresponding carboxylic acids by replacing the ending ‘-ic’ of acid with suffix ‘aldehyde’.
- The IUPAC names of open chain aliphatic aldehydes and ketones are derived from the names of the corresponding alkanes by replacing the ending ‘-e’ with ‘-al’ and ‘-one’ respectively. e.g.,

HCHO,  
Methanal  
(Formaldehyde)

CH<sub>3</sub>—CHO,  
Ethanal  
(Acetaldehyde)

  
Benzencarbaldehyde  
(Benzaldehyde)


  
Butan-2-one

  
Phenylethanone  
(Acetophenone)

  
Diphenylmethanone  
(Benzophenone)


## STRUCTURE

- The C -atom of carbonyl group is  $sp^2$ -hybridised and forms three  $\sigma$  -bonds and one  $\pi$  -bond with O -atom.
- In carbonyl group, carbon and other three atoms lie in the same plane with bond angle of  $120^\circ$ , therefore carbonyl group is planar and  $\pi$ -electron cloud lies above and below of this plane.



## POLAR NATURE OF CARBONYL GROUP


- In carbonyl group,  $>C=O$  bond is stronger, shorter and polarised. As oxygen is more electronegative than carbon, the double bond of carbonyl group is polar and shows dipole moment. Polarisation contributes to the reactivity of aldehydes and ketones.




Dipole moment,  $\mu = 2.3-2.8$  D



## METHODS OF PREPARATION




### Aldehydes and Ketones



### PEEP INTO PREVIOUS YEARS

1. The major product of the following reaction is



(JEE Main 2019)

2. The correct statement regarding a carbonyl compound with a hydrogen atom on its alpha-carbon, is

- a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as carbonylation
- a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as keto-enol tautomerism
- a carbonyl compound with a hydrogen atom on its alpha-carbon never equilibrates with its corresponding enol
- a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as aldehyde-ketone equilibration.

(NEET Phase-I 2016)




## PHYSICAL PROPERTIES

- **Physical state and odour :**
  - Lower members of aldehydes and ketones (upto  $C_{10}$ ) are colourless, volatile liquids except formaldehyde which is gas at ordinary temperature.
  - Higher members of aldehydes and ketones are solids with fruity odour.
  - Lower aldehydes have unpleasant odour but ketones possess pleasant smell.
- **Boiling points :**
  - The boiling points of aldehydes and ketones are higher than hydrocarbons and ethers of comparable molecular masses due to intermolecular attraction (weak dipole-dipole interactions).
  - Their boiling points are lower than those of alcohols of similar molecular masses because these dipole-dipole interactions are weaker than intermolecular hydrogen bonding between alcohol molecules.
  - Among isomeric aldehydes and ketones, ketones have slightly higher boiling points due to the presence of two electron releasing alkyl groups which make carbonyl group more polar.
- **Solubility :**
  - Lower members of aldehydes and ketones (upto  $C_4$ ) are soluble in water due to H-bonding between polar carbonyl group and water. However, solubility decreases with increase in molecular mass.
  - Aromatic aldehydes and ketones are much less soluble than corresponding aliphatic aldehydes and ketones due to larger hydrocarbon part.
  - All carbonyl compounds are fairly soluble in organic solvents.






## CHEMICAL PROPERTIES

- **Nucleophilic addition reactions :**




Carbonyl carbon is electron deficient hence acts as electrophile. Nucleophile attacks on the electrophilic carbon atom of the carbonyl group from a direction perpendicular to the plane of the molecule.



Aldehydes are generally more reactive than ketones towards nucleophilic addition reactions.



- **Nucleophilic addition-elimination reactions :**



### COMIC CAPSULE

I don't trust atoms....





I heard they make up everything.




• **Oxidation and reduction reactions :**

**Oxidation**




• **Haloform reaction :**




(Given by compounds having  $\text{CH}_3\text{CO}$ -group or  $\text{CH}_3\text{CH}(\text{OH})$ -group).


• **Aldol condensation :**




- On heating with dilute acid aldols undergo dehydration to form  $\alpha, \beta$ -unsaturated carbonyl compounds.




- **Cross aldol condensation :** When condensation takes place between two different aldehydes or ketones, or between one aldehyde and one ketone, it is called as cross aldol condensation. If both the reactants contain  $\alpha$ -hydrogen atoms then it gives a mixture of four products.




If only one carbonyl compound contains  $\alpha$ -hydrogen then only one product is formed.



- **Intramolecular aldol condensation :** It takes place in dialdehydes/diketones or compound containing one aldehyde and one ketone group and gives rise to cyclic products.




- **Cannizzaro's reaction :** It is shown by aldehydes which do not have  $\alpha$ -H atom. The aldehydes undergo disproportionation i.e., self oxidation-reduction process.







- **Cross Cannizzaro reaction** : It takes place between two different aldehydes to give all possible products. Cross Cannizzaro reaction of formaldehyde with other aldehydes always gives sodium formate and alcohol of other aldehyde.




- **Intramolecular Cannizzaro reaction** : It is given by dialdehydes having no  $\alpha$ -hydrogen atoms in the presence of concentrated alkali solution.




- **Tischenko's reaction** : All aldehydes (with or without  $\alpha$ -hydrogen atom) in presence of aluminium ethoxide,  $[\text{Al}(\text{OC}_2\text{H}_5)_3]$  undergo simultaneous oxidation (to carboxylic acid) and reduction (to alcohol) to form an ester.



- **Benzoin condensation** : Two molecules of aromatic aldehyde on heating with alcoholic solution of KCN undergo condensation to produce benzoin.




- **Polymerisation** : Lower aldehydes readily undergo polymerisation giving different products under different conditions. Polymerization of acetaldehyde also yields different products under different conditions.




Paraldehyde is used as hypnotic, *i.e.*, sleep inducing agent.


- **Perkin reaction** : It is a condensation reaction in which an aromatic aldehyde condenses with aliphatic acid anhydride in presence of sodium salt of same acid to give a condensate which upon hydrolysis yields  $\alpha,\beta$ -unsaturated acid.




- **Knoevenagel reaction** : It involves condensation between active methylene group and carbonyl groups in the presence of base.




- **Reformatsky reaction** : It involves the reaction of aldehydes and ketones with  $\alpha$ -halogenated ester in presence of Zn to yield  $\beta$ -hydroxy ester.



- **Reaction with ammonia** : Formaldehyde reacts with ammonia to form hexamethylene tetramine. Nitration of hexamethylene tetramine produces RDX, a well-known explosive.



- **Wittig reaction** : It involves the reaction between an aldehyde or ketone and a phosphorus ylide to form alkene.





# CONCEPT MAP

# ESSENTIALS OF INORGANIC CHEMISTRY (PART I)

## General Principles and Processes of Isolation of Elements

### Metallurgy

- Minerals** : Naturally occurring chemical substances in the earth's crust obtainable by mining.
- Ores** : The minerals from which metal can be extracted profitably and conveniently.
- Metallurgy** : The entire scientific and technological process used for isolation of the metal from its ore.

### Concentration of Ore

The whole process of obtaining a pure metal from its ore.

- Crushing and Grinding (Pulverisation)** :  
Lumps of ore  $\xrightarrow{\text{Crushers}}$  Small pieces  $\xrightarrow{\text{Ball mill}}$  Fine powder
- Concentration or Benefication of Ore** : It is the removal of gangue or matrix from the powdered ore. The impurities present in an ore are called gangue or matrix.
- Gravity separation or levigation or hydraulic washing** : For oxide ores, based on difference in the densities of ore (heavier) and gangue particles, using Wilfley tables.
- Froth floatation** : For sulphide ores, based on preferential wetting of ore (by oil) and gangue (by water).
- Magnetic separation** : Based on difference in magnetic properties. (Either ore or the impurities associated with magnetic nature.)
- Leaching** : When ore is soluble in a suitable solvent while impurities are not.  
For Al  $\rightarrow$  Baeyer's process : from bauxite ore  
For Au  $\rightarrow$  Mac-Arthur Forest cyanide process : from argentite ore.

### Thermodynamic Principles of Metallurgy

For a reaction to occur,  $\Delta G$  should be negative. A reaction with  $\Delta G$  positive can be made to occur if it is coupled with another reaction having a large negative  $\Delta G$ , so that the net  $\Delta G$  of both the reactions is positive.

**Ellingham Diagram** : Plots of  $\Delta_r G^\circ$  vs  $T$ . These help in predicting the feasibility of thermal reduction of an ore to metal.

- Slope = Positive, because  $\Delta_r G^\circ$  increases with rise in  $T$ .
- Each curve is a straight line except when phase changes take place ( $s \rightarrow l$ ,  $l \rightarrow g$ ).
- Metal oxide placed higher in the diagram can be reduced by the metal placed lower.

### Metals from Concentrated Ore


- Concentrated Ore** : Conversion of concentrated ore into reducible form (oxidation).
  - Calcination** : Converting concentrated ore into oxide by heating it strongly below its melting point in the absence of air
  - Roasting** : The concentrated ore (usually sulphide) is heated strongly, in the presence of excess of air below its melting point.
- Reduction or conversion of oxides to metals** : The ore obtained after calcination or roasting is reduced to metal and choice of reducing agent depends upon the nature of ore.
- Carbon or carbon monoxide is used for oxides of Fe, Cu, Zn, Mg, Co, etc. and the process is called smelting.
- Electropositive metals like Na, Al, Mg or hydrogen are used for reduction of ores of Mn, Cr, Ti, Mo, W, etc.
- Auto-reduction process is used for ores of Pb, Hg, Cu, etc.
- Electrolytic reduction is used for highly electropositive metals.
- Hydrometallurgy or displacement method is used for Ag, Au, etc.

### Refining


Methods	Metals Purified
Liquation	Used for low melting metals like Sn, Pb, Hg, Bi, etc.
Distillation	Used for volatile metals like Zn, Hg, Cd, etc., or metals containing non-volatile impurities.
Poling	Used for metals which contain impurities of their own oxides e.g., Cu.
Cupellation	Used for metals containing easily oxidisable impurities e.g., Ag containing Pb impurities.
Electrolytic refining	Used for metals like Cu, Ag, Au, Al which get deposited at cathode and impurities get deposited at anode. Solution of a soluble metal salt acts as an electrolyte.
Mond's process	Used for refining of Ni.
Zone refining	Used to produce extremely pure metals (semiconductors) like Si, Ge, Ga, B and In.
van Arkel method	Used for ultra-pure metals like Ti, Zr which are used in space technology.

## The p-Block Elements (Group 15 to 18)


### Group-15 Elements

- General electronic configuration** :  $ns^2 np^3$
- General trends in physical properties** :  

- Moscovium (Mc)** : Atomic no. 115
- Chemical Properties**
  - Hydrides** : Form  $MH_3$  type hydrides.
    - Melting point** :  $PH_3 < AsH_3 < SbH_3 < NH_3$
    - Boiling point** :  $BiH_3 > SbH_3 > NH_3 > AsH_3 > PH_3$
  - Halides** : Form  $MX_3$  and  $MX_5$  types of halides.
    - Lewis acid strength** :  $PCl_3 > AsCl_3 > SbCl_3$  and  $PF_3 > PBr_3 > PI_3$
    - Lewis base strength** :  $NH_3 > NBr_3 > NCl_3 > NF_3$


### Group-17 Elements

- General electronic configuration** :  $ns^2 np^5$
- General trends in chemical properties** :  

- Tennessine (Ts)** : Atomic no. 117
- Chemical Properties**
  - Oxidizing power** :  $F_2 > Cl_2 > Br_2 > I_2$
  - Hydrogen halides** :
    - B.Pt. and M.Pt.** :  $HF > HI > HBr > HCl$
    - Dipole moment and Thermal stability** :  $HF > HCl > HBr > HI$
    - Bond length, acidic strength and reducing nature** :  $HF < HCl < HBr < HI$
  - Oxoacids of halogens** :
    - Acidic strength** :  $HClO < HClO_2 < HClO_3 < HClO_4$

### Group-16 Elements

- General electronic configuration** :  $ns^2 np^4$
- General trends in physical properties** :  

- Livermorium (Lv)** : Atomic no. 116
- Chemical Properties**
  - Hydrides** :  $H_2M$  type, where  $M$  is  $sp^3$  hybridised.
    - M.Pt. and B.Pt.** :  $H_2O > H_2Te > H_2Se > H_2S$
    - Thermal stability** :  $H_2O > H_2S > H_2Se > H_2Te$
    - Volatility** :  $H_2O < H_2Te < H_2Se < H_2S$
    - Reducing power** :  $H_2O < H_2S < H_2Se < H_2Te$
    - Bond angle and dipole moment** :  $H_2Te < H_2Se < H_2S < H_2O$
  - Halides** : Form  $MX_2$ ,  $MX_4$  and  $MX_6$  types of halides.
    - Dihalides** : All elements except selenium form dihalides.
    - Tetrahalides** :  $SF_4$  (gas),  $SeF_4$  (liquid),  $TeF_4$  (solid).  $SF_4$  is readily hydrolysed than  $SeF_4$ .
  - Oxyhalides** : Only S and Se form oxyhalides.
  - Oxides** :  $SO_2$  and  $SeO_2$ ;  $SO_3$ ,  $SeO_3$  and  $TeO_3$  : Acidic
    - $TeO_2$  and  $PO_2$  : Amphoteric

### Group-18 Elements

- General electronic configuration** :  $ns^2 np^6$
- General trends in physical properties** :  

- Oganesson (Og)** : Atomic no. 118
- Chemical Properties**
  - M.Pt. and B.Pt.** are very low.
  - Ionisation enthalpy** :  $He > Ne > Ar > Kr > Xe > Rn$




• **Distinction between aldehydes and ketones :**

Tests	Aldehydes	Ketones
Schiff's reagent	Pink colour	No colour
Fehling's solution	Red precipitate	No precipitate
Tollens' reagent	Silver mirror	No silver mirror
2,4-dinitrophenylhydrazine	Orange-yellow or red well defined crystals with melting points characteristic of individual aldehydes.	Orange-yellow or red well defined crystals with melting points characteristic of individual ketones.

✓ **PEEP INTO PREVIOUS YEARS**

3. Choose the correct option(s) for the following reaction sequence.




Consider Q, R and S as major products.

- (a)   
 (b)   
 (c)   
 (d)


(JEE Advanced 2019)

4. The major product B formed in the following reaction sequence is



(JEE Main Online 2018)

5. Of the following, which is the product formed when cyclohexanone undergoes aldol condensation followed by heating?



(NEET 2017)

6. Reaction of a carbonyl compound with one of the following reagents involves nucleophilic addition followed by elimination of water. The reagent is
- (a) hydrazine in presence of feebly acidic solution   
 (b) hydrocyanic acid   
 (c) sodium hydrogen sulphite   
 (d) a Grignard reagent.

(AIPMT 2015)

🔑 **POINTS FOR EXTRA SCORING**

- $\text{LiAlH}_4$  is more powerful reducing agent than  $\text{NaBH}_4$  as beside aldehydes and ketones it also reduces acids, acid chlorides, esters, amides, anhydrides, nitriles, oximes, alkyl halides, alkyl azides, alkyl tosylates and nitro compounds.  $\text{LiAlH}_4$  and  $\text{NaBH}_4$  do not reduce isolated double bonds.




- Formaldehyde cannot be prepared by Rosenmund's reduction since formyl chloride is unstable at room temperature.
- $\alpha$ -Hydroxy ketones reduce Tollens' reagent, Fehling's and Benedict's solution.
- Haloform reaction can be used to oxidise

$\alpha$ ,  $\beta$ -unsaturated methyl ketones to  $\alpha$ ,  $\beta$ -unsaturated acids without oxidising double bond.

- Oxidation of methyl ketones is governed by Popoff's rule, according to which carbonyl group of the unsymmetrical ketone remains with the smaller alkyl group.


## Carboxylic Acids

- **General Formula** :  $C_nH_{2n}O_2$  having  $-\text{COOH}$  group or  $\text{RCOOH}$  where,  $R=\text{H}$  or alkyl or aryl.
- **Nomenclature** :
  - The common names end with the suffix *-'ic acid'* and have been derived from Latin or Greek names of their natural sources.
  - In the IUPAC system, aliphatic carboxylic acids are named by replacing the ending *-'e'* in the name of the corresponding alkane with *-'oic acid'*. In numbering the carbon chain, the carboxylic carbon is numbered one.




## STRUCTURE

- Carbon-atom of carboxyl group is  $sp^2$ -hybridised and forms one  $\sigma$ -bond with each oxygen atom and one  $\sigma$ -bond with hydrogen or carbon atom depending upon the structure of carboxylic acid. Half-filled  $p$ -orbital of each oxygen atom and unhybridised  $p$ -orbital of carbon-atom lie in the same plane and overlap to form a  $\pi$ -bond which is delocalised between three atoms, one carbon and two oxygen atoms. Thus, the bonds are separated by about  $120^\circ$ .



- Thus, carboxylic acid ( $\text{RCOOH}$ ) can be represented as resonance hybrid of the following structures,



due to which the carboxyl carbon is less electrophilic than carbonyl carbon.

## CLASSIFICATION

- **On the basis of the group to which  $-\text{COOH}$  group is attached :**
  - Aliphatic carboxylic acid :  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$   
(where,  $R = \text{H}$  atom or alkyl group).
  - Aromatic carboxylic acid :  $\text{Ar}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$   
(where,  $\text{Ar} = \text{Aryl}$  group)
- **On the basis of number of  $-\text{COOH}$  groups in their molecule :**

No. of  $-\text{COOH}$  group

  - Monocarboxylic acids : 1
  - Dicarboxylic acids : 2
  - Tricarboxylic acids : 3
- Saturated and unsaturated monocarboxylic acids are also called fatty acids, because higher acids like palmitic acid, steric acid, oleic acid, etc. were obtained by hydrolysis of fats and oils.


## Quotable Quote

Don't read success stories, you will get only message...  
Read failure stories, you will get some ideas to get success...!!

A.P.J. ABDUL KALAM




## Methods of Preparation




## PHYSICAL PROPERTIES

- Acids upto  $\text{C}_{10}$  are liquids with unpleasant odour. The higher members are colourless waxy solids.
- Boiling points** : Boiling points of acids increase regularly with molecular weight and higher than alcohols (of comparable molecular mass) due to formation of dimer through H-bonding



- Solubility** : Acids upto  $\text{C}_4$  are completely soluble in water due to H-bonding. Solubility regularly decreases rapidly due to increase in hydrophobic character of alkyl group.
- Melting points** : The melting point of an acid containing even number of carbon atoms is always higher than the next lower and higher odd number of carbon atoms (alternation effect or oscillation effect) due to effective crystal lattice being symmetrical in nature.




Molecules with even number of C-atoms fit well into the crystal lattice, thus the melting point of these


acids is higher. In higher members, the melting point increases with increase in molecular mass. The aromatic acids have higher melting points than the corresponding aliphatic acids.

## CHEMICAL PROPERTIES


- Reactions involving cleavage of O—H bond** :



- Reactions involving cleavage of C—OH bond** :




- Reactions involving —COOH group** :






- **Hell-Volhard-Zelinsky reaction** : Aliphatic carboxylic acids on reaction with bromine or chlorine in presence of small amount of red phosphorous produces  $\alpha$ -halo acids.



- **Ring substitution in aromatic acids** : Aromatic carboxylic acids undergo electrophilic substitution reactions in which the carboxyl group acts as a *deactivating* and *meta* directing group.




- **Distinction between phenol and carboxylic acid**

Test	Phenol	Carboxylic acid
NaHCO <sub>3</sub> test	No reaction	Brisk effervescence of CO <sub>2</sub> gas
FeCl <sub>3</sub> test	Violet colour	Buff coloured ppt.

### PEEP INTO PREVIOUS YEARS




7. The major product of the following reaction is



- (a)
- (b)
- (c)
- (d)

(JEE Main 2019)

8. The major product of the following reaction is




(NEET 2019)

9. Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their
- formation of intramolecular H-bonding
  - formation of carboxylate ion
  - more extensive association of carboxylic acid *via* van der Waals' forces of attraction
  - formation of intermolecular H-bonding.

(NEET 2018)

10. The correct order of acidity for the following compounds is



- I > II > III > IV
- III > I > II > IV
- III > IV > II > I
- I > III > IV > II


(JEE Advanced 2016)

### POINTS FOR EXTRA SCORING

- Methanoic acid and its sodium salt act as a reducing agent and reduces Tollens' reagent, Fehling's solution and KMnO<sub>4</sub>.


- **Acidic strength of acids** : Benzoic acid ( $K_a = 6.5 \times 10^{-5}$ ) is somewhat stronger than simple aliphatic acids. This is because in benzoic acid the carboxyl group is attached to more electronegative  $sp^2$  hybridised carbon as compared to less electronegative  $sp^3$  hybridised carbon in aliphatic acids.

- Electron withdrawing groups (EWG's) make acids stronger.



- Electron donating groups (EDG's) make acids weaker.






- $-I$  effect increases with increasing number of EWG's.





Strongest acid




Weakest acid

- $-I$  effect decreases with increasing distance from the EWG.



- $\pm R$  effect on aromatic ring decreases or increases acidity.



### Answer Key For Peep Into Previous Years


- |    |     |    |     |    |        |     |     |    |     |    |     |
|----|-----|----|-----|----|--------|-----|-----|----|-----|----|-----|
| 1. | (b) | 2. | (b) | 3. | (a, d) | 4.  | (c) | 5. | (a) | 6. | (a) |
| 7. | (a) | 8. | (c) | 9. | (d)    | 10. | (a) |    |     |    |     |



# WRAP it up!

- The reagent with which both acetaldehyde and acetone react easily is  
(a) Fehling's reagent      (b) Grignard reagent  
(c) Schiff's reagent      (d) Tollens' reagent.





- $\text{C}_6\text{H}_{10}\text{O}_3$  (keto ester)  $\xrightarrow[\Delta]{\text{NaOH} + \text{I}_2}$  yellow ppt. + B  
(A)  
 $B \xrightarrow{\text{H}^+} C \xrightarrow{\Delta} \text{CH}_3\text{COOH}$ .  
Hence, A is



- (c) both of these      (d) none of these.






- 3-Hydroxybutanal is formed when (X) reacts with (Y) in dilute (Z) solution. What are X, Y and Z?  
(a)  $\text{CH}_3\text{CHO}$ ,  $(\text{CH}_3)_2\text{CO}$ , NaOH  
(b)  $\text{CH}_3\text{CHO}$ ,  $\text{CH}_3\text{CHO}$ , NaCl  
(c)  $(\text{CH}_3)_2\text{CO}$ ,  $(\text{CH}_3)_2\text{CO}$ , HCl  
(d)  $\text{CH}_3\text{CHO}$ ,  $\text{CH}_3\text{CHO}$ , NaOH

- Which of the following compounds has wrong IUPAC name?



- $m$ -Chlorobenzaldehyde on reaction with concentrated KOH at room temperature gives  
(a) potassium  $m$ -chlorobenzoate and  $m$ -hydroxy benzaldehyde  
(b)  $m$ -hydroxy benzaldehyde and  $m$ -chlorobenzyl alcohol  
(c)  $m$ -chlorobenzyl alcohol and  $m$ -hydroxybenzyl alcohol  
(d) potassium  $m$ -chlorobenzoate and  $m$ -chlorobenzyl alcohol.










7. Which of the following is used as lachrymator or tear gas?

- (a)  $\text{C}_6\text{H}_5\text{COCl}$  (b)  $\text{C}_6\text{H}_5\text{OC}_6\text{H}_5$   
(c)  $\text{C}_6\text{H}_5\text{COCH}_2\text{Cl}$  (d)  $\text{C}_6\text{H}_5\text{COCH}_3$






8. The end product of the following reaction is



9. Which one of the following is not correct?

- (a)  $\text{>C=O} \xrightarrow{\text{Clemmensen's reduction}} \text{>CH}_2$   
(b)  $\text{>C=O} \xrightarrow{\text{Wolff-Kishner reduction}} \text{>CHOH}$   
(c)  $\text{>COCl} \xrightarrow{\text{Rosenmund's reduction}} \text{>CHO}$   
(d)  $\text{>C}\equiv\text{N} \xrightarrow{\text{Stephen's reduction}} \text{>CHO}$

10. The final product of the following sequence of reactions is



What is B?

- (a)  $\text{CH}_3\text{CH}_2\text{COCl}$  (b)  $\text{CH}_3\text{CH}_2\text{CHO}$   
(c)  $\text{CH}_2=\text{CHCOOH}$  (d)  $\text{ClCH}_2\text{CH}_2\text{COOH}$



12. Aldehyde group can be protected

- (a) by acetal formation against attack by alkaline oxidising agents  
(b) by mercaptal formation against attack by acidic oxidising agents  
(c) by both (a) and (b)  
(d) by none of the above.

13. Benzophenone can be converted into benzene using

- (a) fused alkali  
(b) anhydrous  $\text{AlCl}_3$   
(c) sodium amalgam in water  
(d) acidified dichromate.






14. The number of aldol reaction(s) that occur in the given transformation is



- (a) 1 (b) 2  
(c) 3 (d) 4

15. The general formula  $\text{C}_n\text{H}_{2n}\text{O}_2$  could be for open chain

- (a) diketones (b) carboxylic acids  
(c) diols (d) dialdehydes.



17. Which of the following statements regarding chemical properties of acetophenone are wrong?

- I. It is reduced to methyl phenyl carbinol by sodium and ethanol.  
II. It is oxidised to benzoic acid with acidified  $\text{KMnO}_4$ .




III. It does not undergo electrophilic substitution like nitration at *meta*-position.

IV. It does not undergo iodoform reaction with iodine and alkali.

- (a) I and II (b) II and IV  
(c) III and IV (d) I and II

18. In this reaction,



- (a) 20% *d* + 80% *l*-Isomer  
(b) *d*-Isomer  
(c) *l*-Isomer  
(d) 50% *d* + 50% *l*-Isomer.

19. The increasing order of the rate of HCN addition to compounds I to IV is


- I. HCHO II. CH<sub>3</sub>COCH<sub>3</sub>  
III. PhCOCH<sub>3</sub> IV. PhCOPh  
(a) I < II < III < IV (b) IV < II < III < I  
(c) IV < III < II < I (d) III < IV < II < I

20. The carboxyl functional group (—COOH) is present in

- (a) picric acid (b) barbituric acid  
(c) ascorbic acid (d) aspirin.

## SOLUTIONS



1. (b): Fehling's solution, Schiff's reagent and Tollens' reagent react only with aldehydes, but Grignard reagent reacts with both aldehydes and ketones.



2. (a): A yellow precipitate shows a positive iodoform test. Thus, presence of CH<sub>3</sub>C(=O)—group is confirmed. This keto group is at β-position w.r.t. ester, the β-ketoester can be represented as





Thus, R (by difference) is C<sub>2</sub>H<sub>5</sub>. Thus, β-keto ester is








4. (b)

5. (d): It is an example of Cannizzaro's reaction.



7. (c): Phenacyl chloride is used as 'tear gas'.




12. (c)




c1ccccc1C(=O)c2ccccc2.[K]OH>>c1ccccc1.[K]C(=O)c2ccccc2


Benzophenone + KOH  $\xrightarrow{\text{Fusion}}$  Benzene + Potassium benzoate


$$\begin{array}{c}
 \text{H}_3\text{C}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \xrightarrow[\text{(ii) HCHO (Aldol)}]{\text{(i) OH}^-} \text{HO}-\text{CH}_2-\text{CH}_2-\text{CH}=\text{O} \\
 \downarrow \text{(ii) HCHO (Aldol)} \quad \text{(i) OH}^- \quad \text{(i) OH}^- \quad \text{(ii) HCHO (Aldol)} \\
 \text{HO}-\text{CH}_2-\overset{\text{OH}}{\underset{\text{H}}{\text{C}}}-\text{CHO} \quad \text{HO}-\text{CH}_2-\overset{\text{OH}}{\underset{\text{CH}_2}{\text{C}}}-\text{CHO} \\
 \downarrow \text{(i) OH}^- \quad \text{(ii) HCHO (Cross Cannizzaro's reaction)} \\
 \text{HCOO}^- + \text{HO}-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{OH}
 \end{array}$$


16. (b): 

**18. (d):** Since  $>C=O$  group is planar, nucleophile can attack carbonyl group from either direction and hence a racemic mixture (50% *d* + and 50% *l* - isomer) is obtained in product.

20. (d):   
(Picric acid)

  
(Barbituric acid)

  
(Ascorbic acid)

  
(Aspirin)

For the  
**SCIENTIST** in  
**YOU**

On the artificial leaf, two light absorbers, similar to the molecules in plants that harvest sunlight, are combined with a catalyst made from the naturally abundant element cobalt. When the device is immersed in water, one light absorber uses the catalyst to produce oxygen. The other carries out the chemical reaction that reduces carbon dioxide and water into carbon monoxide and hydrogen, forming the syngas mixture.





# CBSE

## warm-up!

CLASS-XII

Chapterwise practice questions for CBSE Exams as per the latest pattern and marking scheme issued by CBSE for the academic session 2019-20.

### Series 7

### Biomolecules, Polymers, Chemistry in Everyday Life

Time Allowed : 3 hours  
Maximum Marks : 70

#### GENERAL INSTRUCTIONS

- All questions are compulsory.
- Section A: Q.no. 1 to 20 are very short answer questions (objective type) and carry 1 mark each.
- Section B: Q.no. 21 to 27 are short answer questions and carry 2 marks each.
- Section C: Q.no. 28 to 34 are long answer questions and carry 3 marks each.
- Section D: Q.no. 35 to 37 are also long answer questions and carry 5 marks each.
- There is no overall choice. However an internal choice has been provided in two questions of two marks, two questions of three marks and all the three questions of five marks weightage. You have to attempt only one of the choices in such questions.
- Use log tables if necessary, use of calculators is not allowed.

#### SECTION - A

Read the given passage and answer the questions 1 to 5 that follow :

Natural rubber or raw rubber consists of basic material latex which is a dispersion of isoprene. During the treatment this isoprene forms a high molecular weight polymer of isoprene. Majority of the synthetic rubbers are derived from butadiene derivatives. Some important synthetic rubbers are neoprene, Buna-S, Buna-N and thiokol.

- How is natural rubber prepared?
- Define vulcanisation of rubber.
- Name the fillers used in crude rubber and why they are used so?
- What is Buna-S?
- What is Buna-N?

Questions 6 to 10 are one word answers :

- What is the name given to the linkage that holds the amino acids together in proteins?

- To which category, does vulcanised rubber belong?
- Name a watersoluble vitamin which is a powerful antioxidant.
- Name the artificial sweetener which has highest sweetness value in comparison to cane sugar.
- Write the name of one of the common initiators used in free radical addition polymerisation.

Questions 11 to 15 are Multiple choice questions :

- Polymer formation from monomers starts by
  - condensation reaction between monomers
  - coordination reaction between monomers
  - conversion of monomer to monomer ions by protons
  - hydrolysis of monomers.
- Which of the following is a common antacid?
  - NaOH
  - $\text{Ca}(\text{OH})_2$
  - $\text{Mg}(\text{OH})_2$
  - KOH





13. Complete hydrolysis of cellulose gives  
 (a) *D*-fructose (b) *D*-ribose  
 (c) *D*-glucose (d) *L*-glucose.
14. Carbohydrates are stored in human body as the polysaccharide  
 (a) maltose (b) glycogen  
 (c) cellulose (d) amylose.
15. An ester used as medicine is  
 (a) ethyl acetate (b) methyl acetate  
 (c) methyl salicylate (d) ethyl benzoate.

#### Questions 16 to 20 :

- (a) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.  
 (b) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.  
 (c) Assertion is correct, but reason is wrong statement.  
 (d) Assertion is wrong, but reason is correct statement.
16. **Assertion :** Nylon 6, 6 is a thermoplastic polymer.  
**Reason :** It is prepared by condensation polymerisation of hexamethylenediamine and adipic acid.
17. **Assertion :** Acid or enzymatic hydrolysis of sucrose to give an equimolar mixture of *D* (+) glucose and *D* (–) fructose is called inversion.  
**Reason :** Sucrose is the only naturally occurring disaccharide which is reducing sugar.
18. **Assertion :** Penicillin is an antibiotic.  
**Reason :** The drugs which act on the central nervous system and help in reducing anxiety are called antibiotics.
19. **Assertion :** PHBV is an example of biodegradable polymer.  
**Reason :** PHBV is used in speciality packaging, orthopaedic devices, etc.
20. **Assertion :** Ofloxacin is a bactericidal type of antibiotic.  
**Reason :** Ofloxacin has killing effect on microbes.

#### SECTION - B

21. Why is glucose or sucrose soluble in water but cyclohexane or benzene insoluble in water?
22. Write the structures of the monomers of the following polymers:



23. What are natural and synthetic polymers? Give two examples of each type.

OR

How does the presence of double bonds in natural rubber molecules influence their structure and reactivity?

24. (a) A child when dropped his soap in a bathing tub, the soap did not sink but was floating. Suggest the method of manufacturing such soap.  
 (b) Give one example and structure of anionic detergent.
25. Enumerate the reactions of glucose which cannot be explained by its open chain structures.


(Delhi 2014 C, AI 2011, 2010 C)

OR



- (a) Amino acids show amphoteric behaviour. Why?  
 (b) Write one difference between  $\alpha$ -helix and  $\beta$ -pleated structures of proteins. (2018)
26. (a) Sleeping pills are recommended by doctors to the patients suffering from sleeplessness but it is not advisable to take its doses without consultation with the doctor. Why?  
 (b) Which class of drugs is used in sleeping pills?
27. Arrange the following alkenes towards increasing order of reactivity in cationic polymerisation:  
 $\text{H}_2\text{C} = \text{CHCH}_3$ ,  $\text{ClCH} = \text{CH}_2$ ,  
 $\text{H}_2\text{C} = \text{CHC}_6\text{H}_5$ ,  $\text{H}_2\text{C} = \text{CHCOOCH}_3$

#### SECTION - C

28. (a) Answer the following :  
 (i) Can ethylene be polymerised by the cationic process?  
 (ii) Is polyvinyl alcohol soluble in water?  
 (b) Give the structures of each of the products in the following reaction:



29. Write the products in each of the following reactions :






OR

- (a) Differentiate between nucleotide of DNA and RNA.
- (b) The melting point and solubility in water of amino acids are generally higher than that of the corresponding halo acids. Explain.
30. (a) (i) Why is bithional added in soap?  
(ii) Why are soaps biodegradable whereas detergents are non-biodegradable?  
(Delhi 2019)
- (b) Define the following term with a suitable example : Artificial sweeteners.  
(Delhi 2019, AI 2019)

OR

- (a) What are fillers and what role these fillers play in soap?
- (b) How are synthetic detergents better than soaps?
31. (a) Write down the heterogeneous catalyst involved in the polymerisation of ethylene.  
(b) Define the term, 'homopolymerisation' giving an example.
32. State the significance of primary and secondary structures of proteins.
33. (i) Give one example of thermosetting and thermoplastic polymer.  
(ii) Write the monomers of the following polymer :



- (iii) What is the role of sulphur in vulcanization of rubber?
34. (a) What is the basic structural difference between starch and cellulose?  
(b) Give an example of anomers.

#### SECTION - D

35. (a) Aspartame, an artificial sweetener, is a peptide and has the following structure :
- $$\begin{array}{c} \text{CH}_2 - \text{C}_6\text{H}_5 \\ | \\ \text{H}_2\text{N} - \text{CH} - \text{CONH} - \text{CH} - \text{COOCH}_3 \\ | \\ \text{CH}_2 - \text{COOH} \end{array}$$
- (i) Identify the four functional groups.  
(ii) Write its zwitter ion structure.  
(iii) Write the structures of the amino acids obtained from the hydrolysis of aspartame.

- (iv) Which of the two amino acids is more hydrophobic?
- (b) Classify the following into monosaccharides and disaccharides:  
Ribose, 2-deoxyribose, maltose, galactose, fructose and lactose.

OR

An optically active amino acid (A) can exist in three forms depending upon the pH of the medium. If the molecular formula of (A) is  $\text{C}_3\text{H}_7\text{NO}_2$ , write


- (a) the structure of the compound in aqueous medium. What are such ions called?  
(b) in which medium will the cationic form of compound (A) exist?  
(c) in alkaline medium, towards which electrode, the compound (A) migrates in electric field?
36. Answer the following:  
(i) Which material is used as a substitute for wool in making commercial fibres such as orlon or acrilan.  
(ii) What are the conditions required for the preparation of high density polyethene?  
(iii) Name and give the preparation of the polymer used for making unbreakable crockery.

OR

- (a) Account for the fact that radical polymerisation of styrene occurs in a head-to-tail manner.  
(b) Show the head to tail free radical polymerisation of the alkene  $\text{CH}_2 = \text{C}(\text{CH}_3)_2$ .
37. (a) Ranitidine is an antacid. Explain.  
(b) What are food preservatives? Give one example.  
(c) What problem arises in using alitame as artificial sweetener?

OR


- (a) Name the following compound :



- (b) What are antihistamines? Give two examples. Explain how do they act on human body?  
(c) What are the main constituents of dettol?

#### SOLUTIONS

1. Natural rubber is prepared by polymerisation of isoprene.





2. Vulcanisation is a process of treating natural rubber with sulphur or some compound of sulphur under heat as to modify its properties, *i.e.*, to render it non-plastic and to give greater elasticity and ductility.

3. Fillers such as carbon black and zinc oxide are usually added to the crude rubber before vulcanisation in order to improve its wear resistance.

4. Buna-S is a copolymer of butadiene and styrene.

5. Buna-N is a copolymer of butadiene and vinyl cyanide or acrylonitrile.

6. Peptide bond or peptide linkage ( $-\text{CONH}-$ )

7. Elastomer

8. Vitamin C

9. Alitame

10. Benzoyl peroxide

11. (a) : Polymerisation takes place either by condensation or by addition reaction.

12. (c) : Magnesium hydroxide,  $\text{Mg}(\text{OH})_2$  is a common antacid.

13. (c) :  $(\text{C}_6\text{H}_{10}\text{O}_5)_n + n\text{H}_2\text{O} \xrightarrow{\text{H}^+} n\text{C}_6\text{H}_{12}\text{O}_6$   
Cellulose  D-glucose

Cellulose is a straight chain polysaccharide composed of D-glucose units which are joined by  $\beta$ -glycosidic linkages. Hence, cellulose on hydrolysis produces only D-glucose units.

14. (b) : Glycogen is the reserve carbohydrate occurring mainly in the liver.

15. (c) : Oil of winter green (methyl salicylate) is used in medicine.

16. (b)

17. (c) : Sucrose is a non-reducing sugar.

18. (c) : The drugs which act on the central nervous system and help in reducing anxiety are called tranquilizers.

19. (b) : PHBV undergoes bacterial degradation in the environment hence, it is a biodegradable polymer.

20. (a)

21. Sugars like glucose and sucrose have  $-\text{OH}$  groups while cyclohexane and benzene do not. Thus, glucose and sucrose are capable of forming hydrogen bonds with water molecules and hence are soluble in water.

22. (a) Hexamethylenediamine,  $\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}_2$  and adipic acid,  $\text{HOOC}-(\text{CH}_2)_4-\text{COOH}$

(b) Ethylene glycol,  $\text{HOCH}_2-\text{CH}_2\text{OH}$  and

terephthalic acid  $\text{HOOC}-\text{C}_6\text{H}_4-\text{COOH}$

23. **Natural polymers** : Those polymers which are found in plants and animals are called natural polymers *e.g.*, proteins, starch, etc.

**Synthetic polymers** : Polymers which are prepared artificially are called synthetic polymers *e.g.*, plastics like polythene, synthetic fibres such as (nylon 6, 6), etc.

OR

Natural rubber is *cis*-polyisoprene and is obtained by 1, 4-polymerisation of isoprene units. In rubber molecule double bonds are located between  $\text{C}_2$  and  $\text{C}_3$  of each isoprene unit. These *cis*-double bonds do not allow the polymer chains to come closer for effective interactions and hence intermolecular forces are quite weak. Thus, natural rubber *i.e.*, *cis*-polyisoprene has a randomly coiled structure and hence, shows elasticity.

24. (a) Soaps that float in water are made by beating tiny air bubbles into the product before their hardening.

(b)  $\text{CH}_3(\text{CH}_2)_{11}-\text{C}_6\text{H}_4-\text{SO}_3^-\text{Na}^+$

25. The following reactions of D-glucose cannot be explained on the basis of its open chain structure :

(i) D-Glucose does not react with sodium bisulphite ( $\text{NaHSO}_3$ ).

(ii) It does not give 2, 4-DNP test and Schiff's test.

(iii) The pentaacetate of D-glucose does not react with hydroxylamine.

(iv) D-Glucose shows the phenomenon of mutarotation, *i.e.*, when its aqueous solution is kept for sometime its optical activity changes.

(v) On reaction with 1 mole of methanol, it yields two monomethyl derivatives which are known as methyl  $\alpha$ -D-glucoside and methyl- $\beta$ -D-glucoside.

OR

(a) As amino acids have both acidic (carboxyl group) and basic groups (amino group) in the same molecule, they react with both acids and bases. Hence, they show amphoteric behaviour.



(b) In  $\alpha$ -helix structure, intramolecular H-bonding takes place whereas in  $\beta$ -pleated structure, intermolecular H-bonding takes place.

26. (a) If excess of sleeping pills are taken, they may lead to death, therefore they should be taken after consultation with the doctor. Secondly, they are habit forming, therefore, they must be stopped when doctor advises the same.



(b) Tranquilizers are used in sleeping pills.



**27.** Reactivity of an alkene towards cationic polymerisation increases as the stability of intermediate carbocation formed increases. The stability of carbocations increase in the following order.






Therefore, the reactivity of the corresponding alkene towards cationic polymerisation increases as:



**28. (a) (i)** No, a stable carbocation is not formed.

**(ii)** Polyvinyl alcohol can form H-bond with water hence, it is soluble in water.

**(b)**



**OR**

**(a)** Nucleotides are monomeric units of nucleic acids.

A nucleotide is made up of sugar, heterocyclic nitrogenous base and phosphate group.

In case of the DNA nucleotide, the sugar is D-2-deoxyribose whereas in RNA nucleotide, it is D-ribose. They also differ with respect to one base, DNA contains thymine whereas RNA contains uracil.

**(b)** In a solid state, amino acids exist as dipolar ions. Therefore, melting point and solubility in water of amino acids are generally higher than that of the corresponding halo acids.

**30. (a) (i)** Bithional is added in soap to impart antiseptic properties and to reduce the odour produced by bacterial decomposition of organic matter on skin.

**(ii)** Soaps are sodium or potassium salts of long chain fatty acids which can be decomposed by microorganisms

while detergents are sodium salts of long chain alkyl hydrogen sulphate or long chain benzene sulphonic acids which cannot be decomposed by bacteria due to branching.

**(b) Artificial sweeteners :** The substances which provide sweetness to the food without increasing the calories to the body are known as artificial sweetening agents. These are inert and harmless and excreted from the body in urine unchanged. Saccharin and aspartame are artificial sweetening agents.

**OR**


**(a)** Substances which are added to soap to change their properties in order to make them more useful for a particular application, are called fillers. For example,

**(i)** Sodium rosinate, sodium silicate, borax and sodium carbonate are added to laundry soaps to increase their foaming ability.


**(ii)** Glycerol is added to shaving soaps to prevent them from rapid drying.

**(b)** Synthetic detergents are better than soaps because they form foam even in hard water whereas soaps do not. Some of the detergents give foam even in ice cold water.

**31. (a)** The heterogeneous catalyst used for the polymerisation of ethylene is Zeigler Natta catalyst (mixture of trialkyl aluminium,  $\text{R}_3\text{Al}$  and titanium tetra-chloride,  $\text{TiCl}_4$ ).



**(b)** A polymer made by polymerisation of a single monomer is known as homopolymer and the process is known as homopolymerisation *e.g.*, polythene made by polymerisation of ethene molecules.



**32.** The significance of primary structure of proteins is that the amino acid sequence of a protein determines its function and is critical to its biological activity. Even a change of just one amino acid can drastically alter the properties of the entire protein molecule. For example change of glutamic amino acid by valine amino acid in the protein sequence causes sickle cell anaemia.


Secondary structure of proteins provide mechanical movement of muscles, bind substrates to modify them chemically. Antibody proteins bind foreign proteins.



**33. (i)** Thermosetting polymer - Bakelite


Thermoplastic polymer - Polythene.

**(ii)** Melamine-formaldehyde polymer is formed by condensation copolymerisation of melamine and formaldehyde.



**(iii)** Vulcanisation is a process of heating natural rubber with sulphur and an appropriate additive to modify its properties.


It gives greater elasticity and ductility. Sulphur forms cross linked network which gives mechanical strength to the rubber.



**34. (a)** Starch consists of two components, *i.e.*, amylose and amylopectin. Amylose is a linear polymer of  $\alpha$ -D-(+) glucose in which  $\text{C}_1$  of one glucose unit is connected to  $\text{C}_4$  of the other glucose unit through  $\alpha$ -glycosidic linkage. Amylopectin has similar structure but it has branched chain structure. Chain is formed by  $\text{C}_1$ - $\text{C}_4$  glycosidic linkage and branching occurs by  $\text{C}_1$ - $\text{C}_6$  glycosidic linkage.

Cellulose, on the other hand, is a straight chain polysaccharide composed of  $\beta$ -D-glucose and the monomeric glucose units are linked together by  $\beta$ -glycosidic linkage.

**(b)**  $\alpha$ -D-(+)-glucose and  $\beta$ -D-(+)-glucose.



**(i)** In aspartame following four functional groups are present :


(I) ( $-\text{NH}_2$ ) (Amine)

(II) ( $-\text{COOH}$ ) (Carboxylic acid)


(III) ( $-\text{C}(=\text{O})-\text{NH}-$ ) (Amide)

(IV) ( $-\text{C}(=\text{O})-\text{O}-$ ) (Ester)

**(ii)** It's zwitter ion structure is



**(iii)**




**(iv)** Out of the two amino acids [*i.e.* (a) and (b)] formed on hydrolysis, (b) is more hydrophobic because of the presence of non-polar and bulky benzyl group.

**(b) Monosaccharides :** Ribose, 2-deoxyribose, galactose and fructose


**Disaccharides :** Maltose and lactose

**OR**

An optically active amino acid with molecular formula  $\text{C}_3\text{H}_7\text{NO}_2$  is alanine, *i.e.*,



Depending upon the pH of the medium, alanine will exist in the following three forms:



**(a)** In aqueous medium, alanine exists as zwitter ion (I).

**(b)** In acidic medium, it will exist in cationic form (II).

**(c)** In alkaline or basic medium, it exists in anionic form (III) and thus migrates towards anode under the influence of an electric field.

**36. (i)** Polyacrylonitrile, a polymer of acrylonitrile ( $\text{CH}_2=\text{CH}-\text{CN}$ ) is used in the manufacture of acrilan fibres and orlon fibres which are used as a substitute for wool.




(ii) High density polythene is manufactured by heating ethene to 330 K–343 K under the pressure of 6–7 atm in the presence of Ziegler-Natta catalyst. The polythene thus produced practically consists of linear chains of polymer molecules. As those chains pack well, the polymer has high density.


(iii) Melamine and formaldehyde undergo condensation copolymerisation to give melamine-formaldehyde polymer which are used to manufacture the break-resistance crockery under the brand name melmac.

OR

(a) Two radicals are formed depending on the attack of  $OR^\bullet$  (initiator) at the two carbon atoms linked by a double bond in styrene.




Out of these two, the former is more stable being secondary and also stabilized by the phenyl ring, therefore this will control the propagation step. Its formation is also favoured by the fact that attack of  $OR^\bullet$  at the internal carbon atom will be sterically hindered. The stable radical will attack another styrene molecule generating a new radical.



and so on.

The process will continue. The growing polymer chain would thus be joined in a head-to-tail manner.



37. (a) Ranitidine is drug which suppresses the secretion of HCl, therefore, it is called antacid. The

treatment of hyperacidity came through the discovery that histamine stimulates the secretion of pepsin and hydrochloric acid. Ranitidine prevents the interaction of histamine with the receptors (inducing secretion of HCl) present in the stomach wall and thus acts as antacid.

(b) Food preservatives are those chemicals which prevent undesirable changes in flavour, colour and texture of food during storage. They delay these changes and prevent spoilage of food due to microbial growth, e.g., sodium benzoate is commonly used preservative.

(c) Its sweetness cannot be controlled.

OR

(a) The given structure is of bithional.

(b) The hypersensitivity of some people towards dust, pollen grains, cat fur, a particular food, colour or fabric and certain drugs (penicillin, sulpha drugs) is called allergy. Allergy is caused due to release of a substance called histamine by the body, Antihistamines are drugs that either decrease or inhibit the action of histamine in the body, thereby, preventing allergy. Two important antihistamines or antiallergic drugs are brompheniramine and terfenadine. Histamines interact with the binding site of receptors in the body to produce allergy. Antihistamines compete with histamines for these binding sites of the receptors and thereby do not allow histamines to produce allergy.

(c) Chloroxylenol and terpineol.

mtg  
Concerned about your performance in **Class XII Boards?**



#### HIGHLIGHTS

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Paper 2019 Included


For more information or for help in placing your order:



Available at all leading book shops throughout India.  
Visit [www.mtg.in](http://www.mtg.in) for latest offers to buy online!  
Call **0124-6601200** or email: [info@mtg.in](mailto:info@mtg.in)



# MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 120

## Electrochemistry | Chemical Kinetics

Time Taken : 60 Min.


NEET / AIIMS

Only One Option Correct Type

- The electrode potential of a silver electrode dipped in a saturated solution of AgCl in contact with 0.1 M KCl solution at 25°C is  
(Given :  $E^\circ_{\text{Ag}^+/\text{Ag}} = 0.799 \text{ V}$  ;  $K_{sp}$  of AgCl =  $1 \times 10^{-10}$ )  
(a) 0.268 V (b) 0.134 V  
(c) 1.072 V (d) 1.33 V
- For a first order reaction  $A \rightarrow P$ , the temperature ( $T$ ) dependent rate constant ( $k$ ) was found to follow the equation,  $\log k = -(2000)\frac{1}{T} + 6.0$ . The pre-exponential factor  $A$  and the activation energy  $E_a$ , respectively, are  
(a)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $9.2 \text{ kJ mol}^{-1}$   
(b)  $6.0 \text{ s}^{-1}$  and  $16.6 \text{ kJ mol}^{-1}$   
(c)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $16.6 \text{ kJ mol}^{-1}$   
(d)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $38.3 \text{ kJ mol}^{-1}$ .
- A hydrogen electrode placed in a buffer solution of sodium cyanide and HCN in the ratio of  $x : y$  and  $y : x$  has electrode potential value  $a$  and  $b$  volts, respectively, at 25°C. If the difference  $a - b = 35.52 \text{ mV}$ , then the ratio of  $y : x$  is  
(a) 2 (b) 5/2 (c) 3 (d) 2/3
- The rate constant for following reaction :  
 $\text{CH}_3\text{OCH}_3(g) \longrightarrow \text{CH}_4(g) + \text{H}_2(g) + \text{CO}(g)$  at 750 K, is  $6.72 \times 10^{-3} \text{ min}^{-1}$ . Starting with a pressure of 400 mm of Hg at this temperature in a closed container, how many minutes would it take for the pressure in the container to become 760 mm Hg?

- 80 min (b) 40 min  
(c) 89 min (d) 50 min

- The  $E^\circ(\text{V})$  in the given figure is about



- 0.55 (b) 0.61  
(c) 0.78 (d) 0.80
- A drop of solution (volume of 0.05 mL) contains  $3 \times 10^{-6}$  moles of  $\text{H}^+$  ion. If the rate of disappearance of  $\text{H}^+$  ions is  $1 \times 10^7 \text{ mol L}^{-1} \text{ s}^{-1}$ , how long(s) would it take for  $\text{H}^+$  ions in the drop to disappear?

**NEW LAUNCH**

**mtg**

# NEET

## ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus  
24 Mock Tests

Now on your android Smart phones  
with the same login of web portal.


Log on to [test.pcmbtoday.com](http://test.pcmbtoday.com)



- (a)  $6 \times 10^{-6}$  (b)  $6 \times 10^{-9}$   
 (c)  $6 \times 10^{-2}$  (d)  $6 \times 10^{-4}$

7. For the electrode,  $\text{Cu}/\text{Cu}^{2+}$ ,  $\log[\text{Cu}^{2+}]$  (along X-axis) is plotted against  $E_{\text{red}}$  (along Y-axis). The plot obtained is shown in the figure. The electrode potential of the half cell,  $\text{Cu}|\text{Cu}^{2+}$  (0.1 M) will be


- (a)  $-0.34 + \frac{0.0591}{2} \text{ V}$   
 (b)  $0.34 \text{ V}$   
 (c)  $0.34 + \frac{0.0591}{2} \text{ V}$   
 (d)  $-0.34 - \frac{0.0591}{2} \text{ V}$



8. Decomposition of sulphuryl chloride,  $\text{SO}_2\text{Cl}_{2(g)} \longrightarrow \text{SO}_{2(g)} + \text{Cl}_{2(g)}$ . It is found to be a first order reaction. If the initial pressure is  $P_0$  and pressure of the mixture at time  $t$  is ( $P_t$ ), then the rate constant ( $k$ ) would be

- (a)  $k = \frac{2.303}{t} \log \frac{P_0}{2P_0 - P_t}$   
 (b)  $k = \frac{2.303}{t} \log \frac{P_0 - P_t}{P_0}$   
 (c)  $k = \frac{2.303}{t} \log \frac{P_0}{P_0 - P_t}$   
 (d)  $k = \frac{2.303}{t} \log \frac{2P_0}{2P_0 - P_t}$

9. The stability constant of the complex,  $[\text{Zn}(\text{NH}_3)_4]^{2+}$  formed in the following reaction :



[Given :  $E_{\text{Zn}^{2+}/\text{Zn}}^\circ = -0.76 \text{ V}$ ;

$E_{[\text{Zn}(\text{NH}_3)_4]^{2+}/\text{Zn}, 4\text{NH}_3}^\circ = -1.03 \text{ V}$ ]

- (a)  $1.37 \times 10^9$  (b)  $2.35 \times 10^9$   
 (c)  $3.62 \times 10^7$  (d)  $5.05 \times 10^8$
10. The rate constant is numerically equal for three reactions of first, second and third order respectively. Which of the following is true?
- (a) If  $[A] > 1$ ;  $r_3 > r_2 > r_1$   
 (b) If  $[A] = 1$ ;  $r_1 = r_2 = r_3$   
 (c) If  $[A] < 1$ ;  $r_1 > r_2 > r_3$   
 (d) All of these
11. A current of 8.45 A is drawn from a Daniell cell for exactly one hour. If molar mass of Cu and Zn

are 63.5 g/mol and 65.4 g/mol respectively, the loss in mass at anode and gain in mass at cathode respectively are

- (a) 10.0 g, 10.30 g (b) 10.30 g, 10.0 g  
 (c) 16.42 g, 15.35 g (d) 10.73 g, 10.48 g

12. The rate of following reaction :

$\text{H}_2\text{O} + 2\text{S}_2\text{O}_4^{2-}(\text{aq}) \longrightarrow 2\text{HSO}_3^-(\text{aq}) + \text{S}_2\text{O}_3^{2-}(\text{aq})$  studied as follows. In an experiment, the concentration of 0.4 M  $\text{S}_2\text{O}_4^{2-}$  reduces to half in  $3.73 \times 10^6 \text{ s}$ . In another experiment, the concentration 0.25 M  $\text{S}_2\text{O}_4^{2-}$  reduces to half in  $6.0 \times 10^6 \text{ s}$ . The order of the reaction is


- (a) first order (b) second order  
 (c) zero order (d) none of these.

### Assertion & Reason Type

**Directions :** In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.  
 (b) If both assertion and reason are true but reason is not the correct explanation of assertion.  
 (c) If assertion is true but reason is false.  
 (d) If both assertion and reason are false.


NEW LAUNCH



## ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus  
24 Mock Tests for

# JEE Main



Now on your android Smart phones  
with the same login of web portal.

Log on to [test.pcmbtoday.com](http://test.pcmbtoday.com)



13. **Assertion :** In mercury cell, the cell potential is approximately 1.35 V and remains constant during its life.

**Reason :** The overall reaction in mercury cell is represented: as  $\text{Zn(Hg)} + \text{HgO}_{(s)} \rightarrow \text{ZnO}_{(s)} + \text{Hg}_{(l)}$

14. **Assertion :** The kinetic of the reaction,  
 $m\text{A} + n\text{B} + p\text{C} \rightarrow m'\text{X} + n'\text{Y} + p'\text{Z}$

obeys the rate expression as:  $\frac{dx}{dt} = k[\text{A}]^m[\text{B}]^n$ .

**Reason :** The rate of the reaction does not depend upon the concentration of C.

15. **Assertion :** If standard reduction potential for the reaction,  $\text{Ag}^+ + e^- \rightarrow \text{Ag}$  is 0.80 volt, then for the reaction,  $2\text{Ag}^+ + 2e^- \rightarrow 2\text{Ag}$ , it will be 1.60 volt.

**Reason :** If concentration of  $\text{Ag}^+$  ions is doubled, the electrode potential is also doubled.

### JEE MAIN / ADVANCED

#### Only One Option Correct Type

16. For a first order reaction, the ratio of the time taken for  $7/8^{\text{th}}$  of the reaction to complete to that of half of the reaction to complete is  
 (a) 3 : 1 (b) 1 : 3 (c) 2 : 3 (d) 3 : 2
17. A constant current was flown for 1 min through a solution of KI. At the end of experiment, liberated  $\text{I}_2$  consumed 150 mL of 0.01 M solution of  $\text{Na}_2\text{S}_2\text{O}_3$  following the reaction :  
 $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \longrightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$   
 What was the average rate of current flow in ampere?  
 (a) 5.25 A (b) 2.4125 A  
 (c) 3.413 A (d) 4.562 A
18. The Arrhenius equation for the rate constant of decomposition of methyl nitrite and ethyl nitrite are  
 $k_1(\text{s}^{-1}) = 10^{13} \exp\left(\frac{-152300 \text{ J mol}^{-1}}{RT}\right)$   
 and  $k_2(\text{s}^{-1}) = 10^{14} \exp\left(\frac{-157700 \text{ J mol}^{-1}}{RT}\right)$   
 respectively. The temperature at which the rate constants are equal is  
 (a) 282 K (b) 482 K (c) 582 K (d) 382 K
19. Copper sulphate solution (250 mL) was electrolysed using a platinum anode and a copper cathode. A constant current of 2 mA was passed for 16 minutes. It was found that after electrolysis, the absorbance of the solution was reduced to 50%

of its original value. Calculate the concentration of copper sulphate in the solution to begin with.

- (a)  $8.56 \times 10^{-5} \text{ mol L}^{-1}$   
 (b)  $6.52 \times 10^{-5} \text{ mol L}^{-1}$   
 (c)  $7.95 \times 10^{-5} \text{ mol L}^{-1}$   
 (d)  $4.52 \times 10^{-5} \text{ mol L}^{-1}$

#### More than One Options Correct Type

20. The rate constant of a reaction is given by  $k = 2.1 \times 10^{10} \exp(-2700/RT)$ . It means that  
 (a)  $\log k$  versus  $1/T$  will be a straight line with slope =  $\frac{-2700}{2.303 R}$   
 (b)  $\log k$  versus  $1/T$  will be a straight line with intercept on  $\log k$  axis =  $2.1 \times 10^{10}$   
 (c) activation energy ( $E_a$ ) is equal to  $\left(\frac{-2700}{RT}\right)$   
 (d) half life of the reaction decreases with increase of temperature.
21. Consider the following concentration cell :  
 $\text{Zn}_{(s)} | \text{Zn}^{2+} (0.024 \text{ M}) || \text{Zn}^{2+} (0.480 \text{ M}) | \text{Zn}_{(s)}$   
 Which of the following statements is(are) correct?  
 (a) The EMF of the cell at  $25^\circ\text{C}$  is nearly 0.038 V.  
 (b) The EMF of the cell at  $25^\circ\text{C}$  is nearly -0.038 V.  
 (c) If water is added in LHS, so that the  $[\text{Zn}^{2+}]$  is reduced to 0.012 M, the cell voltage increases.  
 (d) If water is added in LHS, so that the  $[\text{Zn}^{2+}]$  is reduced to 0.012 M, the cell voltage remains same.
22. For a first order reaction,  
 (a) The degree of dissociation is equal to  $(1 - e^{-kt})$ .  
 (b) Plot of rate *versus* concentration will be a straight line parallel to concentration axis.  
 (c) The time taken for the completion of 75% reaction is thrice the  $t_{1/2}$  of reaction.  
 (d) The pre-exponential factor in the Arrhenius equation has the dimension of time,  $\text{T}^{-1}$ .
23.  $\text{Pt}_{(s)} | \text{Cl}_2(\text{g}, 1 \text{ atm}) | \text{HCl}(\text{C}_1) || \text{HCl}(\text{C}_2) | \text{Cl}_2(\text{g}, 1 \text{ atm}) | \text{Pt}_{(s)}$ . Mark out the correct statement(s).  
 (a) Cell diagram represents electrode concentration cell.  
 (b) Cell is spontaneous if  $\text{C}_1 > \text{C}_2$ .  
 (c) Cell is spontaneous if  $\text{C}_1 < \text{C}_2$ .  
 (d) Cell diagram represents electrolyte concentration cell.



### Numerical Value Type

24. A current of 1.70 A is passed through 300.0 mL of 0.160 M solution of  $\text{ZnSO}_4$  for 230 sec with a current efficiency of 90%. The molarity (M) of  $\text{Zn}^{2+}$  after the deposition of zinc is (Assume the volume of the solution to remain constant during electrolysis.)
25. The activation energy of a reaction is 9 kcal/mol. The increase in the rate constant when its temperature is raised from 295 to 300 K is  $x\%$ . The value of  $x$  is
26. For a given reaction, energy of activation for forward reaction ( $E_{a_f}$ ) is  $80 \text{ kJ mol}^{-1}$ .  $\Delta H = -40 \text{ kJ mol}^{-1}$  for the reaction. A catalyst lowers  $E_{a_f}$  to  $20 \text{ kJ mol}^{-1}$ . The ratio of energy of activation for reverse reaction before and after addition of catalyst is

### Matrix Match Type

Answer the following questions (27 and 28) by appropriately matching the columns based on the information given in the passage :

At constant temperature and volume, X decomposes as  $2X_{(g)} \rightarrow 3Y_{(g)} + 2Z_{(g)}$ ;  $P_x$  is the partial pressure of X.

S.No.	Time (in minute)	P (in mm of Hg)
1.	0	800
2.	100	400
3.	200	200

Column-I		Column-II	
(P)	Molecularity of reaction	I.	1
(Q)	Order of reaction with respect to X	II.	2
(R)	Time for 75% completion of reaction	III.	$6.93 \times 10^{-3} \text{ min}^{-1}$
(S)	Rate constant	IV.	200 min

27. Which of the following has the correct combination considering column-I and column-II?
- (a)  $P \rightarrow \text{I}$  (b)  $Q \rightarrow \text{III}$   
 (c)  $R \rightarrow \text{IV}$  (d)  $S \rightarrow \text{II}$

28. Which of the following has the correct combination considering column-I and column-II?
- (a)  $P \rightarrow \text{III}$  (b)  $Q \rightarrow \text{I}$   
 (c)  $R \rightarrow \text{II}$  (d)  $S \rightarrow \text{IV}$

Answer the following questions (29 and 30) by appropriately matching the columns based on the information given in the passage :

Arrangement of metals in the order of decreasing tendency of their atoms to give electrons (*i.e.*, undergo oxidation) is known as electrochemical series.

A metal which is placed higher up in the series will displace all metals which are kept below it from solutions of their salts.

- (i)  $\text{Li}^+ + e^- \longrightarrow \text{Li}_{(s)}$ ;  $E^\circ_{\text{Li}^+/\text{Li}} = -3.05 \text{ V}$
- (ii)  $\text{Au}^{3+} + 3e^- \longrightarrow \text{Au}_{(s)}$ ;  $E^\circ_{\text{Au}^{3+}/\text{Au}} = 1.50 \text{ V}$
- (iii)  $\text{F}_{2(g)} + 2e^- \longrightarrow 2\text{F}^-$ ;  $E^\circ_{\text{F}_2/\text{F}^-} = 2.87 \text{ V}$

Column-I		Column-II	
(P)	$\text{F}_2$	I.	Metal is the strongest reducing agent
(Q)	Li	II.	Unreactive metal
(R)	$\text{Li}^+$	III.	Metal ion which is the weakest oxidising agent
(S)	Au	IV.	Non-metal which is best oxidising agent

29. Which of the following has the correct combination considering column-I and column-II?
- (a)  $P \rightarrow \text{III}$  (b)  $Q \rightarrow \text{I}$   
 (c)  $R \rightarrow \text{II}$  (d)  $S \rightarrow \text{IV}$

30. Which of the following has the correct combination considering column-I and column-II?
- (a)  $P \rightarrow \text{IV}$  (b)  $Q \rightarrow \text{II}$   
 (c)  $R \rightarrow \text{I}$  (d)  $S \rightarrow \text{III}$



Keys are published in this issue. Search now! ☺

## SELF CHECK

No. of questions attempted .....  
 No. of questions correct .....  
 Marks scored in percentage .....

### Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.





**C**hemistry Musing was started from August '13 issue of Chemistry Today. The aim of Chemistry Musing is to augment the chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / JIPMER with additional study material. In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today.

The readers who have solved five or more problems may send their solutions. The names of those who send atleast five correct solutions will be published in the next issue. We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.

### JEE MAIN/NEET

1. 1 mol of A in one litre vessel maintained at constant T shows the equilibrium :




If the equilibrium pressure is  $\frac{13}{6}$  times of initial





pressure and  $[C]_{eq} = \frac{4}{9} [A]_{eq}$ , calculate  $K_{C_1}$  and  $K_{C_2}$ .

- (a) 0.11, 0.14 (b) 2.54, 1.46  
(c) 2.96, 0.56 (d) 5.11, 3.21

2. In the given sequence of reactions,




the compound D is

- (a)  (b)   
(c)  (d) 

3. A certain metal (X) is boiled in dilute  $\text{HNO}_3$  to give a salt (Y) and an oxide of nitrogen (Z). An aqueous solution of (Y) with brine solution gives a white precipitate (P). When (Y) is treated with  $\text{Na}_2\text{S}_2\text{O}_3$  solution, a white precipitate (Q) is obtained which on standing turns to a black compound (R). The compound (R) formed is  
(a) PbO (b) PbS (c)  $\text{Ag}_2\text{S}$  (d)  $\text{Ag}_2\text{SO}_3$


4. The gas phase decomposition of dimethyl ether follows first order kinetics.



The reaction is carried out in a container at constant volume and  $500^\circ\text{C}$  and has a half life of 14.5 minutes. Initially only dimethyl ether is present at a pressure of 0.40 atmosphere. The total pressure of the system after 12 minutes is (Assume ideal gas behaviour)

- (a) 0.75 atm (b) 0.90 atm  
(c) 0.56 atm (d) 0.45 atm.

5. Identify (X) in the given reactions.



### GLIMPSE OF NEXT ISSUE...

Monthly Test Drive (XI) :

Monthly Test Drive (XII) :

Brush up for NEET/JEE (XI) :

Brush up for NEET/JEE (XII) :

- Classification of Elements and Periodicity in Properties
- Chemical Bonding
- Surface Chemistry
- General Principles and Processes of Isolation of Elements
- Organic Chemistry - Some Basic Principles and Techniques
- Organic Compound Containing Nitrogen
- Environmental Chemistry



## JEE ADVANCED

6. For the following complexes :

- (1) *trans* -  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$
- (2) *cis* -  $[\text{Co}(\text{NH}_3)_2(\text{en})_2]^{3+}$
- (3) *trans* -  $[\text{Co}(\text{NH}_3)_2(\text{en})_2]^{3+}$
- (4)  $\text{NiCl}_4^{2-}$                       (5)  $\text{TiF}_6^{2-}$




Identify the correct option(s).

- (a) (1) and (2) are optically active, (3) and (5) are optically inactive.
- (b) (2) is optically active, (1), (3) and (4) are optically inactive.
- (c) (4) is coloured, optically active and (5) is colourless, optically inactive.
- (d) (4) is optically inactive, coloured and (5) is optically inactive, colourless.


## COMPREHENSION

On ozonolysis of one mole of compound (M) gives 2 moles of glyoxal and one mole of methyl glyoxal. Compound (M) on reaction with succinic anhydride gives a compound (N) in the presence of anhydrous  $\text{AlCl}_3$ . Compound (N) on heating with Zn-Hg in the presence of HCl gives compound (O). Compound (O) in the presence of conc.  $\text{H}_2\text{SO}_4$  gives compound (P) which in the presence of methyl magnesium chloride followed by hydrolysis gives compound (Q). Compound (Q) on heating in the presence of conc.  $\text{H}_2\text{SO}_4$  gives compound (R) which on heating in the presence of Se gives compound (S).

7. The compound (M) and compound (N) are respectively



8. The compound (S) obtained is



## NUMERICAL VALUE

9. The molar conductances at infinite dilution of  $\text{AgNO}_3$ ,  $\text{NaCl}$ ,  $\text{NaNO}_3$  are 116.5, 110.3 and  $105.2 \text{ mho cm}^2 \text{ mol}^{-1}$  respectively. The electrolytic conductance of  $\text{AgCl}$  in water is  $2.40 \times 10^{-6} \text{ mho/cm}$  and of water used is  $1.16 \times 10^{-6} \text{ mho cm}^{-1}$ . The solubility of  $\text{AgCl}$  is  $x \times 10^{-3} (\text{g litre}^{-1})$ . The value of  $x$  is
10. The total number of cyclic silicates among the following is  
 $\text{Be}_2\text{SiO}_4$  (Phenacite),  $\text{ZrSiO}_4$  (Zircon),  $\text{Mg}_2\text{SiO}_4$  (Forsterite),  $\text{Zn}_4\text{Si}_2\text{O}_7(\text{OH})_2 \cdot \text{H}_2\text{O}$  (Hemimorphite),  $\text{Sc}_2\text{Si}_2\text{O}_7$  (Thortveitite),  $\text{MgSiO}_3$  (Enstatite),  $\text{Ca}_3(\text{Si}_3\text{O}_9)$  (Wollastonite),  $\text{Be}_3\text{Al}_2(\text{SiO}_3)_6$  (Beryl),  $\text{Ca}_2\text{Mg}_5(\text{Si}_4\text{O}_{11})_2$  (Tremolite),  $\text{KAl}_2(\text{OH})_2(\text{AlSi}_3\text{O}_{10})$  (Mica),  $\text{SiO}_2$  (Quartz)

### Solution Senders of Chemistry Musing










#### Set - 76

- Kaushik Dhandekar, Nagpur Maharashtra
- Arkaprov Datta, West Bengal
- Nitun Sarkar, Kolkata

### Solution Senders of Puzzle Corner

- Devjit Acharjee, West Bengal, Kolkata
- Aviral Vishwakarma, New Delhi




- Which of the following is inert towards potassium metal?  
 (a)  (b)   
 (c)  (d) 
- Which of the following has maximum lattice energy?  
 (a) Li<sub>2</sub>O (b) Na<sub>2</sub>O (c) MgO (d) BaO
- Which one of the following is the correct statement?  
 (a) B<sub>2</sub>H<sub>6</sub> · 2NH<sub>3</sub> is known as 'inorganic benzene'.  
 (b) Boric acid is a protonic acid.  
 (c) Beryllium exhibits coordination number of six.  
 (d) Chlorides of both beryllium and aluminium have bridged chloride structures in solid phase.
- $x$  g of Zn metal was mixed with  $2x$  g of iodine to form ZnI<sub>2</sub>. Which of the reactants will remain in excess and what fraction of it remains unreacted? (Zn = 65, I = 127)  
 (a) Zn, 0.75 (b) I<sub>2</sub>, 0.48  
 (c) Zn, 0.48 (d) I<sub>2</sub>, 0.75
- For  $d$ -electron, the orbital angular momentum is  
 (a)  $\frac{\sqrt{6}h}{2\pi}$  (b)  $\frac{\sqrt{2}h}{2\pi}$  (c)  $h/2\pi$  (d)  $2h/\pi$
- The correct order of magnetic moments (spin only values in B.M.) is  
 (a)  $[\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-}$   
 (b)  $[\text{MnCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-}$   
 (c)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-}$   
 (d)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-} > [\text{MnCl}_4]^{2-}$   
 (Atomic nos. : Mn = 25, Fe = 26, Co = 27)
- 0.0833 mol of carbohydrate of empirical formula CH<sub>2</sub>O contain 1 g of hydrogen. The molecular formula of the carbohydrate is  
 (a) C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> (b) C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>  
 (c) C<sub>5</sub>H<sub>10</sub>O<sub>5</sub> (d) C<sub>3</sub>H<sub>4</sub>O<sub>3</sub>
- CsCl crystallises in body-centred cubic lattice. If ' $a$ ' is its edge length then which of the following expressions is correct?  
 (a)  $r_{\text{Cs}^+} + r_{\text{Cl}^-} = \sqrt{3}a$  (b)  $r_{\text{Cs}^+} + r_{\text{Cl}^-} = 3a$   
 (c)  $r_{\text{Cs}^+} + r_{\text{Cl}^-} = \frac{3a}{2}$  (d)  $r_{\text{Cs}^+} + r_{\text{Cl}^-} = \frac{\sqrt{3}}{2}a$
- A closed vessel contains equal number of molecules of N<sub>2</sub> and O<sub>2</sub> at a total pressure of 650 mm of Hg. If N<sub>2</sub> is completely removed, the pressure (mm of Hg) will  
 (a) drop to 650/3 (b) drop to 650/2  
 (c) remain unchanged (d) become 650 × 2
- Which of the following organometallic compound is  $\sigma$  and  $\pi$ -bonded?  
 (a)  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2]$  (b)  $[\text{PtCl}_3(\eta^2\text{-C}_2\text{H}_4)]$   
 (c)  $[\text{Co}(\text{CO})_5\text{NH}_3]^{2+}$  (d)  $\text{Al}(\text{CH}_3)_3$
- CH<sub>3</sub>COOH is neutralised by NaOH. Conductometric titration curve will be  
 (a)   
 (b)   
 (c)   
 (d) 
- Which of the reagents would effect the conversion?  
  
 (a) BH<sub>3</sub>/THF, then H<sub>2</sub>O<sub>2</sub>/OH<sup>-</sup>  
 (b) H<sub>2</sub>O/Hg(OAc)<sub>2</sub>, THF then NaBH<sub>4</sub>/OH<sup>-</sup>  
 (c) H<sub>3</sub>O<sup>+</sup>, Heat  
 (d) None of these.









13. The correct sequence of decrease in the bond angle of the following hydrides is

- (a)  $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3$   
 (b)  $\text{NH}_3 > \text{AsH}_3 > \text{PH}_3 > \text{SbH}_3$   
 (c)  $\text{SbH}_3 > \text{AsH}_3 > \text{PH}_3 > \text{NH}_3$   
 (d)  $\text{PH}_3 > \text{NH}_3 > \text{AsH}_3 > \text{SbH}_3$

14. What are A, B and C in the following sequence of reaction?




- (a)  in all cases  
 (b)  in all cases  
 (c)  in all cases  
 (d) A : , B : ,  
 C : 

15. Which of the following reagents is used to determine the presence of hydroxide groups in reducing sugar?

- (a)  $\text{PCl}_5$  (b)  $\text{HIO}_4$   
 (c)  $(\text{CH}_3\text{CO})_2\text{O}$  (d)  $\text{HI/Red P.}$

16. Consider the following compounds :




The correct order of  $\text{S}_{\text{N}}1$  reactivity is

- (a)  $A > B > C$  (b)  $B > C > A$   
 (c)  $B > A > C$  (d)  $C > B > A$

17. Which series of reactions correctly represents chemical relations related to iron and its compound?

- (a)  $\text{Fe} \xrightarrow{\text{O}_2, \text{heat}} \text{Fe}_3\text{O}_4 \xrightarrow{\text{CO, } 600^\circ\text{C}} \text{FeO} \xrightarrow{\text{CO, } 700^\circ\text{C}} \text{Fe}$   
 (b)  $\text{Fe} \xrightarrow{\text{dil. H}_2\text{SO}_4} \text{FeSO}_4 \xrightarrow{\text{H}_2\text{SO}_4, \text{O}_2} \text{Fe}_2(\text{SO}_4)_3 \xrightarrow{\text{heat}} \text{Fe}$   
 (c)  $\text{Fe} \xrightarrow{\text{O}_2, \text{heat}} \text{FeO} \xrightarrow{\text{dil. H}_2\text{SO}_4} \text{FeSO}_4 \xrightarrow{\text{heat}} \text{Fe}$   
 (d)  $\text{Fe} \xrightarrow{\text{Cl}_2, \text{heat}} \text{FeCl}_3 \xrightarrow{\text{heat, air}} \text{FeCl}_2 \xrightarrow{\text{Zn}} \text{Fe}$


18. The major product expected from the following reaction is



19. If a automobile radiator holds 1.0 kg of water, how many grams of ethylene glycol ( $\text{C}_2\text{H}_6\text{O}_2$ ) must be added to get the freezing point of the solution lowered to  $-2.8^\circ\text{C}$ ? [ $K_f$  for water is  $1.86 \text{ K kg mol}^{-1}$ ]

- (a) 93 g (b) 39 g (c) 27 g (d) 72 g

20. Which of the following structure of  $\text{ClF}_3$  is most stable?



- (d) All the three are of equivalent stability.


#### Numerical Value Type

21. Sea water contains around  $6.5 \times 10^{-3} \text{ g}$  of bromide ions in  $1 \text{ dm}^3$ . If all the bromide ions are converted into  $\text{Br}_2$  how many  $\text{dm}^3$  of sea water are needed to produce 1 kg of bromine?
22. A reaction proceeds five times more at  $60^\circ\text{C}$  as it does at  $30^\circ\text{C}$ . The energy of activation is \_\_\_\_\_.




23. A compound with molecular mass 180 is acylated with  $\text{CH}_3\text{COCl}$  to get a compound with molecular mass 390. The number of amino groups present per molecule of the former compound is \_\_\_\_.
24. 3-Methylpent-2-ene on reaction with  $\text{HBr}$  in presence of  $\text{H}_2\text{O}_2$  forms an addition product. The number of possible stereoisomers for the product is \_\_\_\_.
25. One mole of ethanol is treated with one mole of ethanoic acid at  $25^\circ\text{C}$ . One-fourth of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be \_\_\_\_.

### SOLUTIONS



2. (c): In  $\text{MgO}$ ,  $\text{Mg}^{2+}$  ion is smallest in size and has double the charge in comparison to  $\text{Li}^+$  and  $\text{Na}^+$  ions.

3. (d):



Initial no. of moles  $\frac{x}{65}$   $\frac{2x}{254}$  or  $\frac{x}{127}$

$\text{I}_2$  is in less amount hence  $\text{I}_2$  is the limiting reagent.

$\therefore$  Zn will remain in excess.

$$\text{Number of moles of Zn left unreacted} = \frac{x}{65} - \frac{x}{127}$$

$\therefore$  Fraction of Zn remained unreacted

$$= \frac{\frac{x}{65} - \frac{x}{127}}{\frac{x}{65}} = 0.48$$


5. (a): For  $d$ -electron,  $l = 2$

$$\text{Orbital angular momentum} = \sqrt{l(l+1)} \frac{h}{2\pi}$$


For  $d$ -electron orbital angular momentum

$$= \sqrt{2(2+1)} \frac{h}{2\pi} = \frac{\sqrt{6}h}{2\pi}$$


6. (a):



Number of unpaired electrons = 5



Number of unpaired electrons = 3



Number of unpaired electrons = 0

$$\text{Magnetic moment} = \sqrt{n(n+2)}$$

where  $n$  = number of unpaired electrons.

i.e., greater the number of unpaired electrons, greater will be the paramagnetic character.

7. (a): 0.0833 mol of carbohydrate has 1 g hydrogen.

So, 1 mol of carbohydrate contains  $\frac{1}{0.0833}$  g of hydrogen = 12 g.

$\text{CH}_2\text{O}$  has 2 g of hydrogen.

$$\text{Hence, } n = \frac{12}{2} = 6$$

$\therefore$  Molecular formula of carbohydrate  
 $= (\text{CH}_2\text{O})_6 = \text{C}_6\text{H}_{12}\text{O}_6$

8. (d): In a body-centred cubic (bcc) lattice, oppositely charged ions touch each other along the cross-diagonal of the cube. In case of  $\text{CsCl}$ ,  $2r_{\text{Cs}^+} + 2r_{\text{Cl}^-} = \sqrt{3}a$

$$\text{or, } r_{\text{Cs}^+} + r_{\text{Cl}^-} = \frac{\sqrt{3}}{2}a$$

9. (b): Equal number of molecules means equal number of moles i.e., equal mole fraction.

Since, partial pressure  $\propto$  mole fraction.

$$\therefore p_{\text{N}_2} = p_{\text{O}_2}$$

$$p_{\text{O}_2} = \frac{650}{2} \text{ mm of Hg}$$

### Monthly Test Drive CLASS XII ANSWER KEY


- |           |           |           |             |            |
|-----------|-----------|-----------|-------------|------------|
| 1. (a)    | 2. (d)    | 3. (a)    | 4. (c)      | 5. (b)     |
| 6. (b)    | 7. (a)    | 8. (a)    | 9. (a)      | 10. (d)    |
| 11. (b)   | 12. (b)   | 13. (a)   | 14. (a)     | 15. (d)    |
| 16. (a)   | 17. (b)   | 18. (a)   | 19. (c)     | 20. (a,d)  |
| 21. (a,c) | 22. (a,d) | 23. (c,d) | 24. (0.154) | 25. (28.8) |
| 26. (2.0) | 27. (c)   | 28. (b)   | 29. (b)     | 30. (a)    |




10. (c) : Metal carbonyls are  $\sigma$ - as well as  $\pi$ -bonded.

11. (a) : Initially the conductance of the solution decreases because the addition of NaOH causes not only replacement of  $H^+$  by  $Na^+$  but also suppresses the dissociation of acetic acid but very soon conductance starts increasing with the addition of NaOH due to highly conducting  $OH^-$  ions.

12. (a) :




13. (a) : Bond angle of group 15 element hydrides follows the order :




This is due to decrease in electronegativity difference from N to Sb.

14. (d) :



15. (c) : Acetic anhydride is used to determine the presence of hydroxide groups.

16. (b) :  $S_N1$  reaction rate depends upon the stability of the carbocation, as carbocation formation is the rate determining step. Compound (B), forms a  $2^\circ$  allylic carbocation which is the most stable, the next stable carbocation is formed from (C), it is a  $2^\circ$  carbocation, (A) forms the least stable  $1^\circ$  carbocation, the order of reactivity is thus,



Buy digital copies  
of your favorite  
magazine



Visit Now : [digital.mtg.in](http://digital.mtg.in)

Study Anytime ! Anywhere !




mtg



**17. (a) :** Formation of  $\text{Fe}_3\text{O}_4$  through Fe, corresponds to the combustion of Fe and rest part of the reactions correspond to the production of Fe by reduction of  $\text{Fe}_3\text{O}_4$  in blast furnace.

**18 (d) :**



**19. (a) :**  $K_f = 1.86 \text{ K kg mol}^{-1}$

$$\Delta T_f = 0 - (-2.8) = 2.8^\circ\text{C}$$

Mass of solvent = 1.0 kg

Mass of solute = ?

Molecular mass of solute = 62

$$\Delta T_f = K_f \times m$$

$$m = \frac{\text{Weight of solute}}{\text{Molecular mass of solute}} \times 1000$$


$$m = \frac{\text{Mass of solvent (g)}}{\text{Molecular mass of solute}} \times 1000$$

$$m = \frac{w/62}{1000} \times 1000 = \frac{w}{62}$$

$$\Delta T_f = K_f \times m$$

$$2.8 = 1.86 \times \frac{w}{62} \Rightarrow w = \frac{62 \times 2.8}{1.86} = 93 \text{ g}$$

**20. (c)**



$$6.5 \times 10^{-3} \text{ g Br}^- = 3.25 \times 10^{-3} \text{ g Br}_2 \text{ is in } 1 \text{ dm}^3$$

$$1000 \text{ g Br}_2 \text{ is in } = \frac{1000}{3.25 \times 10^{-3}} = 307.69 \times 10^3 \text{ dm}^3$$

**22. (10.76) :** As  $2.303 \log \frac{k_2}{k_1} = \frac{E_a}{R} \left( \frac{T_2 - T_1}{T_2 T_1} \right)$

Here,  $T_2 = 60 + 273 = 333 \text{ K}$ ,

$T_1 = 30 + 273 = 303 \text{ K}$

$R = 1.987 \times 10^{-3} \text{ kcal}$   $E_a = ?$

As  $r = k[A]^n$  (at a temperature  $T$ )

Hence,  $\frac{r_2}{r_1} = \frac{k_2}{k_1}$  (at temperature  $T_2$  and  $T_1$ )



but  $\frac{r_2}{r_1} = 5$  (Given)

$$\therefore \frac{k_2}{k_1} = 5$$

$$\therefore 2.303 \log_{10} 5 = \frac{E_a}{1.987 \times 10^{-3}} \left[ \frac{333 - 303}{333 \times 303} \right]$$


$$\therefore E_a = 10.76 \text{ kcal mol}^{-1}$$

**23. (5) :** No. of amino groups =  $\frac{390 - 180}{42} = 5$



There are two chiral carbon atoms present in the product. Therefore, total number of stereoisomers are  $= 2^n = 2^2 = 4$

**25. (0.11) :**



Initially	1	1	0	0
At eqm.	$1 - \frac{1}{4}$	$1 - \frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$
	$= 3/4$	$= 3/4$		

$$K_c = \frac{[\text{CH}_3\text{COOC}_2\text{H}_5][\text{H}_2\text{O}]}{[\text{C}_2\text{H}_5\text{OH}][\text{CH}_3\text{COOH}]} = \frac{\left(\frac{1}{4}\right)\left(\frac{1}{4}\right)}{\left(\frac{3}{4}\right)\left(\frac{3}{4}\right)} = 0.11$$



### Monthly Test Drive CLASS XI ANSWER KEY

1. (a)	2. (c)	3. (d)	4. (d)	5. (a)
6. (c)	7. (a)	8. (c)	9. (b)	10. (a)
11. (c)	12. (c)	13. (c)	14. (b)	15. (b)
16. (b)	17. (a)	18. (d)	19. (b)	20. (a,c,d)
21. (a,d)	22. (a,b,c,d)	23. (b,c,d)	24. (4)	
25. (2.64)	26. (20)	27. (a)	28. (c)	
29. (a)	30. (c)			






# CONCEPT BOOSTER



**Dear students!!** Hope you all are doing good. As the season of exams is knocking at your door, now gear up and fasten your seat belt to be safe from anxiety of lacking strategy. This is the high time that you start to revise chapter-wise and more accurately topic-wise to get into the final detailing of the subject. This time revising theory and instantly solving problems will help you a lot. This is a perfect strategy. As you all are my "bestest" friends, so like every year I am there to help you with the proper understanding of every topic with accurate problems. This issue deals with the same. Good luck!!

\*Arunava Sarkar



## Single Option Correct





Identify 'C' in the above sequence of reactions.




Identify the final product.



3. Identify the product.





Identify 'C' in the above sequence of reactions.




\*Institute of Chemistry (IOC)- Asansol, Durgapur, Dhanbad, Burdwan, Kolkata, Jamshedpur, Bokaro, Patna









7. Identify the major product in the following reaction.





8. Arrange the following according to their increasing order of basicity.




Identify the final product.




Identify the final product.



Identify the product.










(d) None of these


### Solutions

1. (c) : In presence of acetic anhydride,  $\text{-NH}_2$  group is prone to be converted to  $\text{-NHCOCH}_3$ .  $\text{NO}_2\text{Cl}$  has polarisation as  $\overset{\delta+}{\text{NO}_2}\overset{\delta-}{\text{Cl}}$ . So,  $\overset{+}{\text{NO}_2}$  will attack the ring activated by both  $\text{-NHCOCH}_3$  and  $\text{-C}_2\text{H}_5$  groups but being more powerful one directive influence will be controlled by  $\text{-NHCOCH}_3$ , hence *para*-position is blocked. So, substitution will surely be at the *ortho*-position.  $\text{NaOH}/\text{CH}_3\text{OH}$  will convert  $\text{-NHCOCH}_3$  back to  $\text{-NH}_2$  as if hydrolysis taking place.




$\text{-NO}_2$  is *meta*-directing. Hence  $\text{-Br}$  will be inserted at the *meta*-position.



3. (b) : It is a mere case of Hoffmann elimination. Moist  $\text{Ag}_2\text{O}$  ( $\text{Ag}_2\text{O} + \text{H}_2\text{O} \rightarrow 2\text{AgOH}$ ) will give  $\ddot{\text{O}}\text{H}$  which will abstract proton from the less hindered site. Therefore, the reaction comes out to be as below :



4. (c) : You know that  $\text{Pb}(\text{OAc})_4$  gives Criegee reaction. So, the first step will be as below along with second step.




Now, in presence of dilute alkali, the product will undergoes internal aldol condensation reaction as below:






6. (c):







$-\text{NH}_2^+$  group is not protonated as the lone pair very much participate in resonance with the benzene ring.

7. (a):



8. (d): (I)



Now, in between (I) and (II), (I) is more basic as in (II), participation of the lone pair for resonance is quite high. In (III), one lone pair very much available. So, the order is,  $\text{II} < \text{I} < \text{III} < \text{IV}$ .



$$\begin{array}{ccc}
 \text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{NH}_2)(\text{NO}_2) & \xrightarrow[0^\circ\text{C}-5^\circ\text{C}]{\text{NaNO}_2, \text{HCl}} & \text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{N}_2^+\text{Cl}^-)(\text{NO}_2) \\
 & & \downarrow \text{HPF}_6 \\
 \text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{F})(\text{NO}_2) & \xleftarrow{\Delta, (-\text{PF}_5)} & \text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{N}_2^+\text{PF}_6^-)(\text{NO}_2)
 \end{array}$$

Electron donating group. This makes the lone pair of nitrogen atom attached with the benzene ring directly more available to react with the outer reagent without intervening into resonance.



# Rank Enhancer

This column is specially designed to make your concepts crystal clear.

## PERIODIC PROPERTIES

The properties which show gradual change when we move from left to right in a period and top to bottom in a group are called periodic properties.

- **Atomic size** : It is defined as distance between centre of nucleus of an atom to its outermost shell of electrons. It is expressed in terms of following three radius :

- Covalent radius, (b) Metallic radius,
- van der Waals' radius

It depends on following factors :

- Number of shells** : Atomic radius increases with increase in number of shells.
- Effective nuclear charge** : Atomic radius decreases with increase in effective nuclear charge.
- Multiplicity of bond** : Atomic radius decreases with increase in multiplicity of bonds.

Example :

Molecule	CH <sub>3</sub> —CH <sub>3</sub>	CH <sub>2</sub> =CH <sub>2</sub>	CH≡CH
Covalent radius of C-atom	77 pm	67 pm	60 pm

### Variation of atomic radii :

- On moving down the group the atomic radius increases due to increase in number of shells.
- On moving left to right in a period, atomic radius decreases due to increase in effective nuclear charge.


### Some Deep Focus Points

- Increase in magnitude of atomic radii is more in the beginning of a group since, difference in energy between energy shells is more.
- Decrease in magnitude of atomic radii is more among representative elements than among transition elements, since electrons are added to penultimate shell.
- Small increase in atomic radii in end of a period is due to the fact that, van der Waals' radius is measured for noble gases.  
van der Waals' radius > covalent radius

- **Ionisation enthalpy** : It is the amount of energy required to remove most loosely bound electron from an isolated gaseous atom.



### Successive ionisation enthalpies :



**Units of I.E.** : 1 eV/atom = 23.06 kcal/mol  
= 96.3 kJ/mol

### Factors affecting I.E. :

- Atomic size :  $I.E. \propto \frac{1}{\text{Atomic size}}$
- Screening effect :  $I.E. \propto \frac{1}{\text{Screening effect}}$
- Effective nuclear charge :  $I.E. \propto \text{nuclear charge}$
- Electronic configuration :  
fully filled > half filled > normal configuration
- Penetration effect :  $I.E. \propto \text{penetration effect}$   
and order of penetration effect is  $s > p > d > f$

### Variation of I.E.

- In a period, as we move from left to right I.E. increases due to decrease in atomic size.
- In a group, as we move down, ionisation energy decreases due to increase in size.

### Some Deep Focus Points

- In second period,  
 $I.E._1 \Rightarrow \text{Li} < \text{Be} > \text{B} < \text{C} < \text{N} > \text{O} < \text{F} < \text{Ne}$   
 $I.E._2 \Rightarrow \text{Li} > \text{Be} < \text{B} > \text{C} < \text{N} < \text{O} > \text{F} < \text{Ne}$
- In third period,  
 $I.E._1 \Rightarrow \text{Na} < \text{Mg} > \text{Al} < \text{Si} < \text{P} > \text{S} < \text{Cl} < \text{Ar}$   
 $I.E._2 \Rightarrow \text{Na} > \text{Mg} < \text{Al} > \text{Si} < \text{P} < \text{S} > \text{Cl} < \text{Ar}$
- Element with lowest I.E. = Cs  
Element with highest I.E. = He
- $Z_{\text{eff}} = Z - \sigma$ , where  
 $Z_{\text{eff}}$  = effective nuclear charge  
 $Z$  = Actual nuclear charge  
 $\sigma$  = Slater's constant (or) screening constant


By K. Vijay Bhasker, Senior faculty at Sri Chaitanya Educational Institution, Visakhapatnam



Electrons in subshells	$\sigma$ per electrons		
	$n^{\text{th}}$ shell	$(n-1)^{\text{th}}$ shell	$(n-2)(n-3)$ ----- etc.
s or p	0.35	0.85	1.00
d or f	0.35	1.00	1.00

### Electron Gain Enthalpy or Electron Affinity

Amount of enthalpy change when an electron is added to an isolated gaseous atom.



### Factors affecting electron affinity :

- Atomic size :  $E.A. \propto \frac{1}{\text{Atomic size}}$
- Effective nuclear charge :  $\Delta_{eg}H \propto Z_{\text{eff}}$
- Stability of electronic configuration : Elements having fully filled (or) half filled configuration have very low (or) zero value of  $\Delta_{eg}H$  and  $\Delta_{eg}H = +ve$ .

### Variation of E.A. :

- Variation in period : As we move from left to right, electron affinity increases as atomic size decreases.
- Variation in group : As we move down, electron affinity decreases.

### Some Deep Focus Points

- Among all elements, Cl has maximum E.A.
- Among metals, gold has high E.A.
- $I.E. \text{ of } X = E.A. \text{ of } X^+$



- Electronegativity** : Measure of tendency of an atom to attract shared pair of electrons towards itself.

### Factors affecting electronegativity :

- Atomic size :  $E.N. \propto \frac{1}{\text{Atomic size}}$
- Nuclear charge :  $E.N. \propto \text{nuclear charge}$
- Screening effect :  $E.N. \propto \frac{1}{\text{Screening effect}}$
- %s character :  $E.N. \propto \% \text{ s character}$

- Electronegativity scales** : Electronegativity is tendency not a physical quantity. Hence, scales are used to compare and for quantitative measurements.

### (1) Pauling scale (Bond energy scale) :

$$\chi_A - \chi_B = 0.1017\sqrt{\Delta} \quad (\text{if } \Delta \text{ in kJ/mol})$$

$$= 0.208\sqrt{\Delta} \quad (\text{if } \Delta \text{ in kcal/mol})$$

$$\text{and } \Delta = E_{A-B} - \frac{1}{2}(E_{A-A} + E_{B-B})$$

**Note :** (i) For reference, take  $\chi_A = 2.1$

(ii) On this scale, F is found to have maximum electronegativity (= 4).

(2) Mulliken scale :  $\chi_A = \frac{I.E. + E.A.}{2}$

$$\chi_A = \frac{I.E. + E.A.}{5.6} \quad (\text{if eV}) = \frac{I.E. + E.A.}{2 \times 62.5} \quad (\text{if kcal/mol})$$

$$= \frac{I.E. + E.A.}{544} \quad (\text{if kJ/mol})$$

Relation between Mulliken and Pauling scale :

$$\Rightarrow \chi_{\text{Pauling}} = \frac{\chi_{\text{Mulliken}}}{2.8}$$

(3) Allred-Rochow scale :  $E.N. = \frac{0.359 \times Z_{\text{eff}}}{r^2} + 0.744$

$r$  = covalent radius,  $Z_{\text{eff}}$  = effective nuclear charge

### Variation of electronegativity :

- As we move from left to right in a period, electronegativity increases.
- As we move down the group, electronegativity decreases.

### Summary of Variation of Periodic Properties

	Periodic property	Down the group	Left to right in period
1.	Atomic size	Increases	Decreases
2.	Ionisation enthalpy	Decreases	Increases
3.	Electron affinity	Decreases	Increases
4.	Electronegativity	Decreases	Increases
5.	Metallic nature	Increases	Decreases
6.	Acidic nature of oxides	Decreases	Increases
7.	Valency	Constant	First increases then decreases
8.	Reducing ability	Increases	Decreases

### QUESTIONS FOR PRACTICE


#### Single Option Correct Type

- If first three successive ionisation energies of an element A are 520, 7297 and 11810 kJ mol<sup>-1</sup> respectively, element belongs to  
(a) group 15      (b) group 18  
(c) group 1      (d) group 2.
- Most basic oxide among the following is  
(a) SiO<sub>2</sub> (b) Na<sub>2</sub>O (c) CO<sub>2</sub> (d) SO<sub>2</sub>
- In inert gases, element with highest electron gain enthalpy is  
(a) He      (b) Ne  
(c) Rn      (d) all have same.



# Precision Revision for JEE is Here!

mtG



## FEATURES:

- 19 years solved papers with detailed solutions
- 10 Model Test Papers
- Chapter-wise indexing of questions

₹625

Now, create your own pre-JEE. Just like pre-boards. With previous years' papers and model test papers for JEE Advanced, complete with detailed solutions, identify your areas of weakness and work on addressing them in time. Multiple test papers ensure you do your dry runs again and again, till such time you feel confident of taking on the best. For it will indeed be the best you compete with in JEE Advanced. So what are you waiting for? **Order MTG's JEE Advanced Explorer today.**



Scan now with your  
smartphone or tablet  
Application to read  
QR codes required

Available at all leading book shops throughout India. To buy online visit [www.mtg.in](http://www.mtg.in).

For more information or for help in placing your order, call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)



4. Mass fraction of hydrogen in hydride of 14<sup>th</sup> group element is 0.125. Then the formula of hydride is  
(a) CH<sub>4</sub> (b) GeH<sub>4</sub> (c) SiH<sub>4</sub> (d) SnH<sub>4</sub>
5. Which of the following electronic configuration corresponds to maximum electron affinity?  
(a) 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>5</sup> (b) 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>5</sup>  
(c) 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>3</sup> (d) 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>3</sup>
6. The ionic radii (in Å) of N<sup>3-</sup>, O<sup>2-</sup> and F<sup>-</sup> respectively are  
(a) 1.71, 1.40, 1.36 (b) 1.71, 1.36, 1.40  
(c) 1.36, 1.40, 1.71 (d) 1.36, 1.71, 1.40
7. Electronegativity of Be is nearly equal to  
(a) B (b) Mg (c) Al (d) Si
8. If configurations of elements P, Q, R, S are  
P → 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> Q → 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>1</sup>  
R → 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>4</sup> S → 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>5</sup>  
Correct order of electron accepting tendency is  
(a) P < R < Q < S (b) P < Q < R < S  
(c) S < R < Q < P (d) S < Q < R < P
9. If bond energies of A—A, H—H and H—A are 38, 104 and 138 kcal/mol respectively, electronegativity of A is  
(a) 3.5 (b) 3.8 (c) 3.0 (d) 1.7
10. Which of the following is not a fixed quantity for an element?  
(a) I.E.<sub>1</sub> (b) E.A.  
(c) Electronegativity (d) I.E.<sub>2</sub>

### Matching List Type

11. Match the following :

	Electronic configuration		Electron gain enthalpy
(A)	1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup> 3s <sup>1</sup>	(P)	-328
(B)	1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>5</sup>	(Q)	-141
(C)	1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>4</sup>	(R)	-53
(D)	1s <sup>2</sup> 2s <sup>2</sup> 2p <sup>6</sup>	(S)	+48

12. Match the following :

	Element		Slater's constant
(A)	Be	(P)	4.15
(B)	Na	(Q)	2.05
(C)	Ne	(R)	1.70
(D)	Li	(S)	8.80

### Hints & Solutions

1. (c) : As 1<sup>st</sup> ionisation energy of A is very low, it can't belong to 15, 16 groups and there is sudden increase in I.E.<sub>2</sub>. Hence, it belongs to group 1.

2. (b) : Metal oxides are basic and non-metal oxides are acidic. In given oxides, Na being most metallic, Na<sub>2</sub>O is most basic.

3. (b) : He < Rn < Xe < Kr = Ar << Ne  
(+48) (+68) (+77) (+96) (+96) (+116)

4. (c) : For 14<sup>th</sup> group, valency will be 4 hence general formula will be AH<sub>4</sub>. Let mass of A = m.

$$\text{Then mass fraction of H} = \frac{(4 \times 1)}{(4 \times 1) + (m \times 1)} = 0.125$$

$$\therefore m = 28$$

Atomic mass of Si = 28  $\therefore$  Formula is SiH<sub>4</sub>.

5. (b) : In options (c) and (d), elements belong to 15<sup>th</sup> group whereas in options (a) and (b) elements belong to 17<sup>th</sup> group. E.A. of 17<sup>th</sup> group > 15<sup>th</sup> group and option (b) corresponds to Cl, (a) corresponds to F.

As (E.A.)<sub>Cl</sub> > (E.A.)<sub>F</sub> therefore, option (b) is correct.

6. (a) :

	N <sup>3-</sup>	O <sup>2-</sup>	F <sup>-</sup>
Electrons	10	10	10
Protons	7	8	9

No. of protons per 10e<sup>-</sup> increases

$\therefore$  Effective nuclear charge increases and ionic radius decreases.

Be	B
Mg	Al

7. (c) :

Be shows diagonal relationship with Al.

8. (b) : P < Q < R < S  
↓ ↓ ↓ ↓  
Inert gas Metal Chalcogen Halogen

9. (b) :  $\Delta = 138 - \frac{1}{2}(38 + 104) = 67$

$$\chi_A - \chi_H = 0.208 \sqrt{67}$$

$$\chi_A - 2.1 = 0.208 \sqrt{67} \Rightarrow \chi_A = 3.8$$

10. (c) : Electronegativity is a tendency but not quantity.

11. A → R ; B → P ; C → Q ; D → S

Halogens have more -ve EGE. Metals have less -ve EGE and noble gases has +ve EGE.

12. A → Q ; B → S ; C → P ; D → R

$$\text{Be} \Rightarrow 1s^2 2s^2 = (2 \times 0.85) + 1 \times 0.35 = 2.05$$

$$\text{Na} \Rightarrow 1s^2 2s^2 2p^6 3s^1 = (2 \times 1) + (0.85 \times 8) = 8.80$$

$$\text{Ne} \Rightarrow 1s^2 2s^2 2p^6 = (2 \times 0.85) + (7 \times 0.35) = 4.15$$

$$\text{Li} = 1s^2 2s^1 = (0 \times 0.35) + (2 \times 0.85) = 1.70$$







# How to overcome exam stress and anxiety

Exams whether of school, Board or of any competition like NEET, AIIMS, JEE, etc. have grown into a do or die pressure-giving event which makes students go through a lot of stress and anxiety. This can cause major mental health issues if allowed to develop. The fear of these exams casts dire consequences on students. The time which the students could have used in having a focused preparation goes wasted in worrying for the exams. Hence, the solution to this problems lies in systematic planning followed by actually making the plan work by following it.

**Here are some tips to get rid of exam fear and anxiety :**

➤ **Plan an early revision**

By now, students might have completed their curriculum and started revision. Practice, practice and do more practice by solving as much as questions and problems related to your subjects.

➤ **Plan a productive timetable**

To start for a result-oriented preparation, formulation of a productive timetable is extremely necessary. Hence, the subjects that the students consider lesser prepared, would rule their timetable. The students give more time to these disciplines. Draw plan for each day and each week. During their daily study, things like inclusion of breaks and their duration, addressing their other important jobs etc, will need students regulate them on daily basis. Hence, students must prepare themselves to deal with these each day.

Also, students may try sticking to a schedule of fixing a few days or a week for each of their subjects. The approach will likely help them to complete their preparation in a timely manner.

➤ **Consider incorporation of more subjects into daily studies**

Preparing for a single subject for hours and hours may get quite monotonous for students. Hence, they can consider extending their devotion to a few more subjects in a single day.

➤ **Prepare notes**

Preparing notes while studying will help the students immensely. These

would provide them the facility of getting support as and when they desire.

Try to make notes of important facts, formulae, unit conversions in the form of table, flowcharts, etc.

Moreover, students can also use a highlighting pen for attracting their attention to words/phrases or (parts of) sentences they need to prepare, more quickly.

➤ **Practice solutions to question papers of previous years**

Following such an exercise will provide the students the opportunities for refining their concepts. It will indicate their stand for the upcoming exams. The questions and problems in these papers will provide them almost the same environment as the actual exams will provide them. This is because they will reflect the same pattern as will be there in the exams. Furthermore, students can turn their study-room into a real exam-room if they stipulate their preparation with the general 3-hour limit. Such an arrangement would really provide them almost real outcomes regarding their preparation.

MTG Previous Years' Questions Bank and Mock Test Papers series of various exams will be very helpful for students.

➤ **Schedule regular breaks, have a balanced meal and sleep well**

The students must consider the size, number and the ways to spend their breaks, when giving breaks to their preparation. They can fully avail this liberty here! They must only be mindful of the fact that they need to maintain a good balance between their breaks and study session. The duration of the sleep may depend on requirements from student to student. Still, a 7-hour slumber would be an ideal one. Students must take a healthy meal filled with all the necessary five nutrients- protein, carbohydrates, fats, minerals, and vitamins.

Students must not compromise with the outcomes of their efforts on preparation, come what may! So, keep the stress aside, and get down to work!

**All the Best!** 🍀



Best **Tips/Informations** that help you to crack **2020 Exams**





Visit **blog.pcmbtoday.com**



# CHEMISTRY MUSING

## SOLUTION SET 76

1. (d): Reactions take place at electrodes are :



For the deposition of 197 g (1 mole) of  $\text{Au} = 3F$  of charge is required thus, for the deposition of 78 g of

$$\text{Au, charge required} = \frac{3}{197} \times 78 = 1.19 F$$

$$= 1.19 \times 96500 \text{ coulombs}$$

$$\text{From } Q = I \times t$$

$$t = \frac{Q}{I} = \frac{1.19 \times 96500}{6} = 19139.16 \text{ sec}$$

By 1 F charge, 35.5 g of  $\text{Cl}_2$  gas is formed, thus from 1.19 F, mass of chlorine gas formed =  $35.5 \times 1.19 = 42.24 \text{ g}$

2. (a): Mass of electron =  $\frac{1}{1835} \times \text{Mass of proton}$   
K.E. of electron =  $16E$

$$\text{Mass of proton} = m_p$$

$$\text{K.E. of proton} = 4E$$

$$\text{Mass of } \alpha\text{-particle} = 4m_p$$



$$\text{K.E. of } \alpha\text{-particle} = E$$

$$\lambda_e = \frac{h}{\sqrt{2 \left( \frac{m_p}{1835} \right) \times 16E}} = \frac{h\sqrt{1835}}{4\sqrt{2 \times m_p E}}$$

$$\lambda_p = \frac{h}{\sqrt{2 \times m_p \times 4E}} = \frac{h}{2\sqrt{2m_p E}}$$

$$\lambda_\alpha = \frac{h}{\sqrt{2 \times 4m_p \times E}} = \frac{h}{2\sqrt{2m_p E}}$$

$$\lambda_e > \lambda_p = \lambda_\alpha$$




5. (d): Heat of hydrogenation depends upon number of  $\pi$ -bonds. When number of  $\pi$ -bonds are same then stability of the compound depends upon the number of alkyl group attached to doubly bonded carbon atom. Greater the number of alkyl group more stable is the alkene.

$$\text{Heat of hydrogenation} \propto \frac{1}{\text{Stability of compound}}$$

6. (a,d): Assume 1 F charge is passed through  $\text{FeSO}_4$ ,  $\text{Fe}_2(\text{SO}_4)_3$  and  $\text{Fe}(\text{NO}_3)_3$ .

	$\text{FeSO}_4$	$\text{Fe}_2(\text{SO}_4)_3$	$\text{Fe}(\text{NO}_3)_3$
Amount of Fe deposited	$\frac{56}{2}$	$\frac{56}{3}$	$\frac{56}{3}$

7. (a): The reaction for the formation of pentane is



For *n*-pentane :

$$\Delta_r H^\circ = \Delta_f H^\circ (\textit{n}\text{-pentane}) - 5\Delta_f H^\circ (\text{C}_{(s)}) - 6\Delta_f H^\circ (\text{H}_{2(g)})$$

$$\text{or } \Delta_r H^\circ = -200 \text{ kJ/mol}$$

For *n*-pentane :  $\Delta_r S^\circ$

$$= \Delta_f S^\circ (\textit{n}\text{-pentane}) - 5\Delta_f S^\circ (\text{C}_{(s)}) - 6\Delta_f S^\circ (\text{H}_{2(g)})$$

$$\text{or } \Delta_r S^\circ = 360 - 5 \times 6 - 6 \times 130 = 360 - 810$$

$$= -450 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\therefore \Delta_r G^\circ = \Delta_r H^\circ - T\Delta_r S^\circ = -200 + 400 \times 0.450 = -20 \text{ kJ/mol}$$

8. (d):  $5\text{C}_{(s)} + 6\text{H}_{2(g)} \longrightarrow \text{C}_5\text{H}_{12(g)}$   
(Mixture of *n*-pentane, *iso*-pentane and *neo*-pentane)

$$\therefore \text{For } \textit{n}\text{-pentane, } \ln K_{eq1} = -\frac{\Delta_r G^\circ}{RT} = \frac{20}{RT}$$

$$\text{or } K_{eq1} = e^{20/RT}$$

$$\text{For } \textit{iso}\text{-pentane, } K_{eq2} = e^{\frac{22}{RT}} \text{ and}$$

$$\text{For } \textit{neo}\text{-pentane, } K_{eq3} = e^{\frac{26}{RT}}$$

$$\therefore K_{eq1} = \frac{p_n}{p_{\text{H}_2}^6} \therefore p_n = K_{eq1} \times p_{\text{H}_2}^6$$

$$\text{and } p_{\text{iso}} = K_{eq2} \times p_{\text{H}_2}^6 \text{ and } p_{\text{neo}} = K_{eq3} \times p_{\text{H}_2}^6$$

$\therefore$  At equilibrium,

$$x_{\text{neo}} = \frac{K_{eq3}}{K_{eq3} + K_{eq2} + K_{eq1}} = \frac{1}{1 + \frac{K_{eq2}}{K_{eq3}} + \frac{K_{eq1}}{K_{eq3}}}$$



9. (6) : Mass of  $\text{Fe}_2\text{O}_3$  in the given sample  $= \frac{55.2}{100} \times 1$   
 $= 0.552 \text{ g}$

Number of moles of  $\text{Fe}_2\text{O}_3 = \frac{0.552}{159.8} = 3.454 \times 10^{-3}$   
 (Mol. wt. of  $\text{Fe}_2\text{O}_3 = 159.8$ )

1 mole of  $\text{Fe}_2\text{O}_3$  will give 2 moles of  $\text{Fe}^{3+}$   
 $= 2 \times 3.454 \times 10^{-3} \text{ moles of } \text{Fe}^{3+}$   
 $= 6.908 \times 10^{-3} \text{ mol} = 6.908 \text{ mmoles}$

$\text{Fe}^{3+}$  is reduced to  $\text{Fe}^{2+}$  by zinc dust.

Since the conversion of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$  or  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  involves only 1 electron change, so the molecular weight is same as equivalent weight.

Amount of  $\text{Fe}^{2+}$  ion in 100 mL of solution = 6.908 meq.


Volume of oxidant used for 100 mL of  $\text{Fe}^{2+}$  solution  
 $= 17 \times 4 \text{ mL} = 68 \text{ mL}$

Amount of oxidant used =  $68 \times 0.0167 \text{ mmoles}$   
 $= 1.1356 \text{ mmoles}$

Let, the number of electrons taken up by oxidant =  $n$

Number of meq of oxidant used =  $1.1356 \times n$

Thus,  $1.1356 \times n = 6.908$  or  $n = \frac{6.908}{1.1356} \approx 6$



Since pressure after reaction is half of the initial pressure so after reaction gaseous moles,  $x = \frac{1}{2}$

i.e.  $\frac{1}{2}$  mole of  $B$  is formed.

Now the number of moles of oxygen atoms before and after reaction = 2 moles

$\Rightarrow 0.5 \text{ moles of } B \text{ contains} = 2 \text{ moles of O atoms}$

$\Rightarrow 1 \text{ mole of } B \text{ contains} = 4 \text{ moles of O atoms}$

1 molecule of  $B$  contains = 4 atoms of oxygen



## More than half of South Asian youth are not on track to have the education and skills necessary for employment in 2030: UNICEF

An estimated 54 per cent of South Asian youth leave school without the necessary skills to get a decent job in the next decade, according to data produced by the Global Business Coalition for Education (GBC-Education), the Education Commission, and UNICEF. According to the data, South Asia lags behind several other regions in preparing the next generation of young people with the skills they will need for 21st century work. Projections place South Asia well below the global average. This builds on the estimates UNICEF produced with the Commission last year on the projected learning outcomes in 2030 for all South Asian countries.

"Every day, nearly 100,000 young South Asians – a large sports stadium of young people – enter the labour market, almost half of them not on track to find 21st century jobs," said Henrietta Fore, UNICEF Executive Director. "South Asia is at a critical juncture, with a limited window during which it can reap significant demographic dividends from its talented and capable youth. Get it right, and millions could be lifted out of poverty. Fail to do so, and economic growth will falter, youth despair will rise, and further talent will be lost to other regions." With almost half of its population of 1.8 billion below the age of 24, led by India, Pakistan and Bangladesh, South Asia will have the largest youth labour force in the world until 2040. This offers the region the potential to drive vibrant and productive economies. If strong investments in skills development are made, the region is poised to maintain strong economic growth as well as an expansion of opportunities in the education and skills sectors in the coming decades.

However, a recent UNICEF 'Voices of Youth' survey conducted among 32,000 young people in South Asia reveals concerns among these under 24-year-olds about how well they are being prepared for the modern economy. According to the poll, many young people in South Asia feel their education systems are outdated and do not prepare them for employment. They cite lack of work experience (26%), inadequate support services to improve employability (23% received no support and most received limited and not comprehensive support), and bribery demands/discriminatory and unfair hiring practices (44%) as key barriers to finding employment even after they have graduated.

"This is a crisis," said Justin Van Fleet, Executive Director of the Global Business Coalition for Education. "Addressing the youth skills gap in South Asia requires government investments, commitments from the business community, contributions from civil society, and the perspective of young people to best equip the next generation to successfully enter the rapidly changing job market." A separate new report commissioned by UNICEF identified major obstacles to addressing the youth skills gap in the region. These include the low quality of education and suboptimal vocational training which do not give students the desired skill levels the labour market demands, among others. Compiled by Ernst & Young India, this report identifies a total of 30 already-in-operation solutions to the youth skills crisis.

### Children in South Asia on Track to Complete Secondary School & Learn Basic Skills by 2030\*

Based on current trends

Million	Today	2030 Projection	School-Age Children in 2030
Afghanistan	†	†	13.9 Million
Bangladesh	26%	55%	37.8 Million
Bhutan	47%	81%	0.2 Million
India	19%	47%	309.6 Million
Maldives	16%	46%	0.09 Million
Nepal	13%	46%	7.1 Million
Pakistan	18%	40%	64.3 Million
Sri Lanka	61%	68%	3.8 Million

† Afghanistan is not included due to lack of recent learning assessment data at the secondary level.

\* These estimates were generated based on a 2019 update of the Education Commission's original 2016 projections model for the Learning Generation report. Most recent national learning assessment data used for each country as follows: BCSE 2015 for Bhutan, GCE O Levels 2016 for Sri Lanka, LASI 2015 for Bangladesh, NAT 2016 for Pakistan, NCERT 2017 for India, Nepal country assessment 2017 for Nepal, O Level Exam 2016 for Maldives. Afghanistan is not included due to lack of recent learning assessment data at the secondary level.



**mtg**

Now, savings of up to  
**₹920\*** with MTG's  
 magazine  
 subscription plans!

\*On cover price of ₹40/-

**Our new offers are here!**

Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to [www.mtg.in](http://www.mtg.in) now to subscribe online.

**For JEE  
 (Main &  
 Advanced),  
 NEET, AIIMS  
 & JIPMER**

### About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines (Physics For You, Chemistry Today, Mathematics Today & Biology Today) ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?



**Over 1.2 Cr readers. Since 1982.**

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team comprising experts and members from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement – confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET/AIIMS toppers
- Bonus: Exposure to competition at a global level, with questions from International Olympiads & Contests

**Lifetime Subscription Plan for teachers, and special schemes and offers available for libraries and coaching institutes. SMS MTG to 8800255334 to learn more.**

## SUBSCRIPTION FORM

Confirm your choice by placing ☒ tick-marks in relevant boxes.

<b>Plan 1:</b> Individual magazines P, C, M, B	<input type="checkbox"/> Physics <input type="checkbox"/> Chemistry <input type="checkbox"/> Mathematics <input type="checkbox"/> Biology	<b>27 months</b> <input type="checkbox"/> ₹850 (save ₹230)	<b>15 months</b> <input type="checkbox"/> ₹500 (save ₹100)	<b>9 months</b> <input type="checkbox"/> ₹300 (save ₹60)
<b>Plan 2:</b> Combo of 3	<input type="checkbox"/> PCM <input type="checkbox"/> PCB	<input type="checkbox"/> ₹2500 (save ₹740)	<input type="checkbox"/> ₹1400 (save ₹400)	<input type="checkbox"/> ₹900 (save ₹180)
<b>Plan 3:</b> PCMB Combo		<input type="checkbox"/> ₹3400 (save ₹920)	<input type="checkbox"/> ₹1900 (save ₹500)	<input type="checkbox"/> ₹1200 (save ₹240)
<b>Courier Charges</b> <small>Add to your subscription amount for quicker &amp; reliable delivery</small>		<input type="checkbox"/> ₹600	<input type="checkbox"/> ₹450	<input type="checkbox"/> ₹240

**Recommended by (Optional)**

\_\_\_\_\_  
Name of your teacher

Teacher's Mobile #

Note: Magazines are despatched by Book-Post on 4<sup>th</sup> of every month (each magazine separately).

Name: \_\_\_\_\_

Complete Postal Address: \_\_\_\_\_

Pin Code

Mobile #


Other Phone # 0

Email \_\_\_\_\_

Enclose Demand Draft favouring  
**MTG Learning Media (P) Ltd.**, payable at New Delhi.  
 Mail this Subscription Form to Subscription Dept.,  
**MTG Learning Media (P) Ltd.**, Plot 99, Sector 44, Gurugram - 122 003 (HR).

E-mail [subscription@mtg.in](mailto:subscription@mtg.in). Visit [www.mtg.in](http://www.mtg.in) to subscribe online. Call (0)1800-10-38673 for more info.  
 Get digital editions of MTG Magazines on <http://digital.mtg.in/>





# Trust MTG for getting it right, year after year

Over the last 5 years, MTG has averaged a hit rate of 70% when it comes to curating the right content for your NEET preparation. Which means approx. 3 out of 4 questions in NEET were either exactly the same as, or similar to, questions in MTG's NEET books. The credit for this mind-blowing feat goes to MTG's skilled and experienced editorial team which painstakingly goes through volumes of NCERT subject matter that forms the basis for NEET, to create superior and relevant study material that has a high chance of success for its users. Proof lies in the pudding, right!

## MTG's best-selling NEET books include



Scan to buy on [mtg.in](http://mtg.in)

To find out which MTG NEET book is best-suited to your needs,  
call our NEET helpline toll-free at **1800-10-38673** today.  
Or email [info@mtg.in](mailto:info@mtg.in) now. Visit [bit.ly/mtg-neet](http://bit.ly/mtg-neet) to buy online.





**mtg**



Scan to buy on [Amazon.in](http://Amazon.in)



# Presenting India's No.1 NEET Guides



MTG's Complete NEET Guides are India's best selling PMT books!! Rich in theoretical knowledge with a vast question bank comprising a wide variety of problems and exercises, these guidebooks ensure students are ready to compete in the toughest of medical entrance tests. 100% NCERT based, the guidebooks have been updated to match the syllabus and the exam pattern for medical entrance exams. No wonder these guidebooks emerged as the bestsellers in a short period of time.

## HIGHLIGHTS:

- 100% NCERT based
- Comprehensive Chapterwise theory complemented with concept maps, flowcharts and easy-to-understand illustrations
- Last 10 years' questions (2010-2019) of AIPMT/NEET
- Chapterwise Topicwise MCQs with detailed explanations and solutions
- More than 80% same or similar MCQs in NEET are from MTG NEET Books



Scan now with your  
smartphone or tablet\*



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or e-mail: info@mtg.in

\*Application to read QR codes required

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!





# FULLY LOADED & COMPLETELY UPDATED


mtg

MTG's BITSAT Power Guide is not only the most exhaustive prep-tool, but also the only book available at present, updated as per the latest BITSAT syllabus for students aspiring for top rank in BITSAT 2020.



Attempt all questions from this book on the Web + Mobile for **free**

(For details see inside the book)



₹ 850/-

Get MTG's BITSAT Power Guide today for a real-world feel of BITSAT. Find out what's different about the BITSAT test, including its pattern of examination and key success factors. Be it with chapterwise MCQs or model test papers, check how good your chances are for glory in BITSAT 2020.

## FEATURES

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapterwise MCQs in each section for practice
- Past chapterwise BITSAT Qs (2019-2011)
- 10 Model Test Papers with detailed solutions
- Attempt all questions & tests from this book online, for free with #L

Visit [www.MTG.in](http://www.MTG.in) to buy online. Or visit a leading bookseller near you.  
For more information, email [info@mtg.in](mailto:info@mtg.in) or call 1800-10-38673 (toll-free).



# CBSE CHAMPION Chapterwise -Topicwise Solved Papers



CBSE CHAMPION Chapterwise-Topicwise Solved Papers Series contains topicwise questions and solutions asked over last decade in CBSE-Board examination.

Questions are supported with topicwise graphical analysis of previous years CBSE Board questions as well as comprehensive and lucid theory. The questions in each topic have been arranged in descending order as per the marks category. Questions from Delhi, All India, Foreign and Compartment papers are included. This ensures that all types of questions that are necessary for Board exam preparation have been covered.

Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme. CBSE sample paper and practice papers are also supplemented.

Examination papers for Class-10 and 12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.



**SRM**INSTITUTE OF SCIENCE & TECHNOLOGY  
(Deemed to be University u/s 3 of UGC Act, 1956)[www.srmist.edu.in](http://www.srmist.edu.in)KATTANKULATHUR | RAMAPURAM  
VADAPALANI | DELHI - NCR, GHAZIABAD**SRM**UNIVERSITY  
DELHI-NCR, SONEPAT[www.srmuniversity.ac.in](http://www.srmuniversity.ac.in)DELHI - NCR SONEPAT,  
HARYANA**SRM**UNIVERSITY  
Sikkim[www.srmus.ac.in](http://www.srmus.ac.in)

GANGTOK, SIKKIM

**SRM**UNIVERSITY AP  
Andhra Pradesh[www.srmap.edu.in](http://www.srmap.edu.in)

AMARAVATI, AP



# LAUNCH INTO A BRIGHT FUTURE

## SRMJEEE 2020

**SRM Joint Entrance Examination - B.Tech Admissions Open**  
**12<sup>th</sup> April - 20<sup>th</sup> April 2020 [Computer based test]**To apply online and other details visit [www.srmist.edu.in](http://www.srmist.edu.in)For eligibility and fee details, visit the respective websites  
Top 3000 SRMJEEE rank holders will get scholarship**LAST DATE FOR SUBMITTING ONLINE APPLICATION: 30<sup>th</sup> MARCH 2020**  
**ONLINE APPLICATION FEE: ₹1100/-**

### LEAD WITH SRMIST

- Multicultural Learning Environment
- Students from all States of India
- International Students from 64 Countries
- 165+ MOUs with 33 countries
- 300+ students from SRM go to our Semester Abroad Programme (SAP) every year
- SRM Innovation & Incubation Centre (SIIC) funded by National Science and Technology Entrepreneurship Development, DST, Govt. of India
- 2 years + 2 years transfer programmes with our global partners

**Admissions Helpline: 044 2745 5510 ; 044 4743 7500****Email: [admissions.india@srmist.edu.in](mailto:admissions.india@srmist.edu.in)**

SRM University Sikkim offers only Science and Humanities

SRMUniversityOfficial
 SRM\_Univ
 srmuniversityofficial
 SRM IST Chennai

### NUMBERS SPEAK FOR THEMSELVES

**4749**

offers from 4 Top IT firms

**1612+**

Super Dream &amp; Dream offers

The success story of SRM 2020 batch placement continues...

- TCS, Infosys, Cognizant & Wipro make the highest number of offers Pan India
- More than 7000 offers till date and still counting

- IDEAL (Interdisciplinary Experiential Active Learning) equips you with Dual Specialisation • World-class faculty and Infrastructure • International Exposure

**IDEAL**  
INTER DISCIPLINARY EXPERIENTIAL ACTIVE LEARNING

### NATIONAL RECOGNITION - SRMIST

- NAAC "A++" grade • Category I University by UGC
- Ranked 32nd under University Category by NIRF • 12B status by UGC
- ARIIA Ranked Third under the Category of Private or Self financed Institutions

### GLOBAL RECOGNITION - SRMIST

- Quacquarelli Symonds (QS) also awarded SRM 5 stars for Teaching, Employability and Inclusiveness
- QS I-GUAGE Diamond rated Institute



Offline application forms can be obtained:

- From the designated branches of Post Offices on cash payment of ₹1100 (list available on website)
- From SRM campuses by handing over a DD for ₹1100 in favour of SRMIST payable at Chennai





# Excellent Performance

## —by Aakashians in—

### NEET-UG | AIIMS | JEE (Main) | JEE (Advanced)

#### • RESULTS 2019 •

## 80081

69826 Classroom + 10255 Distance & Digital  
**NEET-UG 2019**

8 in Top 10 | 15 in Top 20 | 31 in Top 50 AIR

Our Top Rankers in General Category



**1**  
AIR  
Nalin Khandelwal  
Distance



**2**  
AIR  
Bhavik Bansal  
Classroom



**3**  
AIR  
Akshat Kaushik  
Classroom



**4**  
AIR  
Swastik Bhatia  
Classroom



**5**  
AIR  
Anant Jain  
Classroom

and many more...

## 688

576 Classroom + 112 Distance & Digital  
**AIIMS 2019**

6 in Top 10 | 11 in Top 20 | 25 in Top 50 AIR

Our Top Rankers in General Category



**1**  
AIR  
Bhavik Bansal  
Classroom



**2**  
AIR  
Vishwa Vadodaria  
Distance



**3**  
AIR  
Akshat Kaushik  
Classroom



**4**  
AIR  
Chetanya Mittal  
Distance



**6**  
AIR  
Arunangshu B.  
Distance

and many more...

## 7879

7250 Classroom + 629 Distance & Digital  
**JEE (Main) 2019**

2 in Top 10 | 10 in Top 50 | 16 in Top 100 AIR

Our Top Rankers from Classroom Programs in Open Category



**16**  
AIR  
Dravya Marwaha



**26**  
AIR  
Prateek Chaudhury



**39**  
AIR  
Sarvagya Jain



**56**  
AIR  
Samanyu Mahajan



**122**  
AIR  
Uday Singla

and many more...

## 1633

1441 Classroom + 192 Distance & Digital  
**JEE (Advanced) 2019**

13 in Top 100 | 23 in Top 200 AIR

Our Top Rankers from Classroom Programs in Open Category



**51**  
AIR  
Sayantan Dhar



**58**  
AIR  
Sibasis Nayak



**83**  
AIR  
Prateek Chaudhury



**107**  
AIR  
Tushar Singla



**126**  
AIR  
Dravya Marwaha

and many more...

### CLASSROOM COURSES

#### 1, 2, 3<sup>rd</sup> & 4<sup>th</sup> YEAR INTEGRATED COURSES

JEE (Main & Advanced) &  
Other Engg. Ent. Exams | NEET & AIIMS

For Class IX, X, XI & XII Studying Students

#### POWER STEP COURSE (Repeater Course)

JEE (Main & Advanced) &  
Other Engineering Entrance Exams

For Class XII Passed Students

#### REGULAR COURSE (Repeater Course)

JEE (Main) & Other Engineering  
Entrance Exams | NEET & AIIMS

For Class XII Passed Students

\*For details & availability of 3 & 4 Year Courses, contact the nearest branch/centre of Aakash.

Though every care has been taken to publish the result correctly, yet the institute shall not be responsible for error, if any.

### DIGITAL / DISTANCE LEARNING PROGRAMS

for NEET & AIIMS / JEE (Main & Advanced) - Class XI & XII Studying / Passed Students | For School / Board Exams / NTSE & Olympiads (Class VIII / IX / X)



**Aakash**  
iTutor

RECORDED VIDEO LECTURES



**Aakash**  
Live

LIVE ONLINE CLASSES



**Aakash**  
Distance Learning Program  
Medical | IIT-JEE | Foundations

PRINTED STUDY MATERIAL

For Digital / Distance Learning Programs Call: 8800012992 or Give a Missed Call: 9599698683



**SCAN FOR  
NEAREST  
CENTRE &  
CONTACT DETAILS**

**TOLL-FREE 1800-102-2727**

**Give a missed call 9599280605**



# Aakash

**Medical | IIT-JEE | Foundations**

(Divisions of Aakash Educational Services Limited)

Registered Office: Aakash Tower, 8, Pusa Road, New Delhi-110005. Ph.: (011) 47623456 | E-mail: reach.us@aesl.in | SUNDAY OPEN

www.aakash.ac.in | instagram.com/aakasheducation | facebook.com/aakasheducation | twitter.com/aakash\_twitted | youtube.com/AakashEducation | aakash.ac.in/blogs